This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISI	HED U	INDER THE PATENT COOPERATION TREATY (PCT)
(51) International Patent Classification 7:		(11) International Publication Number: WO 00/23111
A61K 39/395, 48/00, C12P 19/34, C12Q 1/68, G01N 33/53, 33/574, 33/546, 33/567	A1	(43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/US (22) International Filing Date: 19 October 1999 (CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
(30) Priority Data: 60/104,737 19 October 1998 (19.10.98) (71) Applicant (for all designated States except US): DIA LLC [US/US]; 3303 Octavius Drive, Santa Clara, (US).	ADEXU	
(72) Inventors; and (75) Inventors/Applicants (for US only): SALCEDA [AR/US]; 4118 Crescendo Avenue, San Jose, C (US). RECIPON, Herve [FR/US]; 85 Fortuna Ave Francisco, CA 94115 (US). CAFFERKEY, Robert Apartment #218, 350 Elan Village Lane, San . 95134 (US).	A 9512 nue, S [IE/US	66
(74) Agents: LICATA, Jane, Massey et al.; Law Office Massey Licata, 66 E. Main Street, Marlton, NJ 080		
	G, STA	GING, IMAGING AND TREATING PROSTATE CANCER
(57) Abstract		GING, IMAGING AND TREATING PROSTATE CANCER , diagnosing, monitoring, staging, prognosticating, imaging and treating
(57) Abstract The present invention provides new methods for de		
(57) Abstract The present invention provides new methods for de		
(57) Abstract The present invention provides new methods for de		
(57) Abstract The present invention provides new methods for de	etecting	
(57) Abstract The present invention provides new methods for de	etecting	diagnosing, monitoring, staging, prognosticating, imaging and treating
(57) Abstract The present invention provides new methods for de	etecting	diagnosing, monitoring, staging, prognosticating, imaging and treating

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BC	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	ΙE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	- Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PΤ	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD OF DIAGNOSING,

MONITORING, STAGING, IMAGING AND TREATING PROSTATE CANCER

FIELD OF THE INVENTION

This invention relates, in part, to newly developed 5 assays for detecting, diagnosing, monitoring, staging, prognosticating, imaging and treating cancers, particularly prostate cancer.

BACKGROUND OF THE INVENTION

Cancer of the prostate is the most prevalent malignancy in adult males, excluding skin cancer, and is an increasingly prevalent health problem in the United States. In 1996, it was estimated that 41,400 deaths would result from this disease in the United States alone, indicating that prostate cancer is second only to lung cancer as the most common cause of death in the same population. If diagnosed and treated early, when the cancer is still confined to the prostate, the chances of cure is significantly higher.

Treatment decisions for an individual are linked to the stage of prostate cancer present in that individual. A common classification of the spread of prostate cancer was developed by the American Urological Association (AUA). The AUA system divides prostate tumors into four stages, A to D. Stage A, microscopic cancer within prostate, is further subdivided into stages A1 and A2. Sub-stage A1 is a well-differentiated cancer confined to one site within the prostate. Treatment is generally observation, radical prostatectomy, or radiation. Sub-stage A2 is a moderately to poorly differentiated cancer at multiple sites within the prostate. Treatment is radical prostatectomy or radiation. Stage B, palpable lump within the prostate, is also further subdivided into sub-stages B1 and B2. In sub-stage B1, the cancer forms a small nodule in one

WO 00/23111

- 2 -

PCT/US99/24331

lobe of the prostate. In sub-stage B2, the cancer forms large or multiple nodules, or occurs in both lobes of the prostate. Treatment for sub-stages B1 and B2 is either radical prostatectomy or radiation. Stage C is a large cancer mass 5 involving most or all of the prostate and is also further subdivided into two sub-stages. In sub-stage C1, the cancer forms a continuous mass that may have extended beyond the prostate. In sub-stage C2, the cancer forms a continuous mass that invades the surrounding tissue. Treatment for both these 10 sub-stages is radiation with or without drugs to address the cancer. The fourth stage, Stage D is metastatic cancer and is also subdivided into two sub-stages. In sub-stage D1, the cancer appears in the lymph nodes of the pelvis. In sub-stage D2, the cancer involves tissues beyond lymph nodes. Treatment 15 for both of these sub-stages is systemic drugs to address the cancer as well as pain.

However, current prostate cancer staging methods are limited. As many as 50% of prostate cancers initially staged as A2, B, or C are actually stage D, metastatic. Discovery of metastasis is significant because patients with metastatic cancers have a poorer prognosis and require significantly different therapy than those with localized cancers. The five year survival rates for patients with localized and metastatic prostate cancers are 93% and 29%, respectively.

Accordingly, there is a great need for more sensitive and accurate methods for the staging of a cancer in a human to determine whether or not such cancer has metastasized and for monitoring the progress of a cancer in a human which has not metastasized for the onset of metastasis.

It has now been found that a number of proteins in the public domain are useful as diagnostic markers for prostate cancer. These diagnostic markers are referred to herein as cancer specific genes or CSGs and include, but are not limited to: Prol09 which is a human zinc- α 2-glycoprotein (Freje et al. Genomics 1993 18(3):575-587); Prol12 which is a human

- 3 -

cysteine-rich protein with a zinc-finger motif (Liebhaber et al. Nucleic Acid Research 1990 18(13):3871-3879; MD9514772 and WO9845436); Proll1 which prostate-specific is a transglutaminase (Dubbink et al. Genomics 1998 51(3):434-444); 5 Prol15 which is a novel serine protease with transmembrane, LDLR, and SRCR domains and maps to 21q22.3 (Paoloni-Giacobino et al. Genomics 1997 44(3):309-320; WO9837418 and WO987093); Prol10 which is a human breast carcinoma fatty acid synthase (U.S. Patent 5,665,874 and WO9403599); Prol13 which is a 10 homeobox gene, HOXB13 (Steinicki et al. J. Invest. Dermatol. 1998 111:57-63); Pro114 which is a human tetraspan NET-1 (WO9839446); and Proll8 which is a human JM27 protein (WO9845435). ESTs for these CSGs are set forth in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13 and 15 while the full length contigs for 15 these CSGs are set forth in SEQ ID NO:2, 4, 6, 8, 10, 12, 14 and 16, respectively. Additional CSGs for use in the present invention are depicted herein in SEQ ID NO: 17, 18, 19 and 20.

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging, prognosticating, 20 imaging and treating prostate cancer via the cancer specific genes referred to herein as CSGs. For purposes of the present invention, CSG refers, among other things, to native protein expressed by the gene comprising a polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 25 16, 17, 18, 19 or 20. By "CSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1-20, but which still encode the same protein. alternative, what is meant by CSG as used herein, means the .30 native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, levels of the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 35 20, or levels of a polynucleotide which is capable of

hybridizing under stringent conditions to the antisense sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, \clubsuit 0, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of CSG in the patient versus the normal human control is associated with prostate cancer.

Further provided is a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which is not known to have metastasized by identifying a human patient suspected of having prostate cancer that has metastasized; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in CSG levels in the patient versus the normal human control is associated with prostate cancer which has metastasized.

- 5 -

Also provided by the invention is a method of staging prostate cancer in a human which has such cancer by identifying a human patient having such cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

Further provided is a method of monitoring prostate cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided is a method of monitoring the change in stage of prostate cancer in a human having such cancer by looking at levels of CSG in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

- 6 -

Further provided are methods of designing therapeutic agents targeted to a CSG for use in *maging and treating prostate cancer. For example, in one embodiment, therapeutic agents such as antibodies targeted against CSG or 5 fragments of such antibodies can be used to detect or image localization of CSG in a patient for the purpose of detecting or diagnosing a disease or condition. Such antibodies can be polyclonal, monoclonal, or omniclonal or prepared by molecular biology techniques. The term "antibody", as used herein and 10 throughout the instant specification is also meant to include aptamers and single-stranded oligonucleotides such as those derived from an in vitro evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable labels including, 15 but not limited to, radioisotopes and paramagnetic metals. Therapeutics agents such as antibodies or fragments thereof can also be used in the treatment of diseases characterized by expression of CSG. In these applications, the antibody can be used without or with derivatization to a cytotoxic agent 20 such as a radioisotope, enzyme, toxin, drug or a prodrug.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging and prognosticating cancers

- 7 -

by comparing levels of CSG in a human patient with those of CSG in a normal human control. For purposes of the present invention, what is meant be CSG levels is, among other things, native protein expressed the gene by comprising a 5 polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. By "CSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1-20, but which still encode the 10 same protein. The native protein being detected, may be whole, a breakdown product, a complex of molecules or chemically modified. In the alternative, what is meant by CSG as used herein, means the native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 15 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, levels of the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, or levels of a polynucleotide which is capable of hybridizing under stringent conditions to the 20 antisense sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. Such levels are preferably determined in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in 25 accordance with the invention for diagnosing overexpression of CSG protein compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of prostate cancer.

All the methods of the present invention may optionally include determining the levels of other cancer markers as well as CSG. Other cancer markers, in addition to CSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

- 8 -

Diagnostic Assays

The present invention provides methods for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of CSG in the patient versus the normal human control is associated with the presence of prostate cancer.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a human cancer patient suspected of having prostate cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art.

In the present invention, determining the presence of CSG levels in cells, tissues or bodily fluid, is particularly useful for discriminating between prostate cancer which has not metastasized and prostate cancer which has metastasized. Existing techniques have difficulty discriminating between prostate cancer which has metastasized and prostate cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, the cancer marker levels measured in such cells, tissues or bodily fluid is CSG, and are compared with levels of CSG in preferably the same cells, tissue or bodily fluid type of a normal human control. That

- 9 -

is, if the cancer marker being observed is just CSG in serum, this level is preferably compared with the level of CSG in serum of a normal human control. An increase in the CSG in the patient versus the normal human control is associated with prostate cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal patient.

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may preferably also include samples from a human patient that is determined by reliable methods to have prostate cancer which has not metastasized.

20 Staging

The invention also provides a method of staging prostate cancer in a human patient. The method comprises identifying a human patient having such cancer and analyzing cells, tissues or bodily fluid from such human patient for CSG. The CSG levels determined in the patient are then compared with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG (but still increased over true normal levels) is associated with a cancer which is regressing or in remission.

Monitoring

Further provided is a method of monitoring prostate 35 cancer in a human patient having such cancer for the onset of

- 10 -

WO 00/23111 PCT/US99/24331

metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; and comparing the CSG levels determined in the human patient with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which has metastasized. In this method, normal human control samples may also include prior patient samples.

Further provided by this invention is a method of monitoring the change in stage of prostate cancer in a human patient having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; and comparing the CSG levels determined in the human patient with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease in the levels of CSG is associated with a cancer which is regressing in stage or in remission. In this method, normal human control samples may also include prior patient samples.

25 Monitoring a patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression (including protein levels), such as CSG of the present invention, in a sample derived from a patient are well known to those of skill in the art. Such assay methods include, without limitation, radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry

- 11 -

assays, in situ hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based approaches such as mass spectrometry or protein interaction profiling. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific 10 to CSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to CSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to CSG is incubated on a solid support, e.g. a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific 20 protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time CSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to CSG and linked to a detectable 25 reagent such as horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to CSG. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added -30 to the dish. Immobilized peroxidase, linked to CSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of CSG protein present in the sample. Quantitative results typically are obtained by reference to a standard 35 curve.

WO 00/23111

- 12 -

A competition assay can also be employed wherein antibodies specific to CSG are attached to a solid support and labeled CSG and a sample derived from the host are passed over the solid support. The amount of label detected which is attached to the solid support can be correlated to a quantity of CSG in the sample.

PCT/US99/24331

Nucleic acid methods can also be used to detect CSG mRNA as a marker for prostate cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain 10 reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reversetranscriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population 15 in a complex mixture of thousands of other mRNA species. RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR RT-PCR can thus reveal by amplification the 20 presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e. gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the CSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the CSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest. Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but

- 13 -

not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by in vitro transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

Of the proteomic approaches, 2D electrophoresis is a 10 technique well known to those in the art. Isolation of individual proteins from a sample such as serum accomplished using sequential separation of proteins by different characteristics usually on polyacrylamide gels. First, proteins are separated by size using an electric The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since 20 no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative 25 abundance of a given protein and the identity of the proteins in the sample.

The above tests can be carried out on samples derived from a variety of cells, bodily fluids and/or tissue extracts such as homogenates or solubilized tissue obtained from a patient. Tissue extracts are obtained routinely from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva or any other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum or any derivative of blood.

- 14 -

In Vivo Targeting of CSGs

Identification of these CSGs is also useful in the rational design of new therapeutics for imaging and treating cancers, and in particular prostate cancer. For example, in 5 one embodiment, antibodies which specifically bind to CSG can be raised and used in vivo in patients suspected of suffering from prostate cancer. Antibodies which specifically bind a CSG can be injected into a patient suspected of having prostate cancer for diagnostic and/or therapeutic purposes. 10 The preparation and use of antibodies for in vivo diagnosis is well known in the art. For example, antibody-chelators labeled with Indium-111 have been described for use in the radioimmunoscintographic imaging of carcinoembryonic antigen expressing tumors (Sumerdon et al. Nucl. Med. Biol. 1990 15 17:247-254). In particular, these antibody-chelators have been used in detecting tumors in patients suspected of having recurrent colorectal cancer (Griffin et al. J. Clin. Onc. 1991 9:631-640). Antibodies with paramagnetic ions as labels for use in magnetic resonance imaging have also been described 20 (Lauffer, R.B. Magnetic Resonance in Medicine 1991 22:339-Antibodies directed against CSG can be used in a similar manner. Labeled antibodies which specifically bind CSG can be injected into patients suspected of having prostate cancer for the purpose of diagnosing or staging of the disease 25 status of the patient. The label used will be selected in accordance with the imaging modality to be used. For example, radioactive labels such as Indium-111, Technetium-99m or Iodine-131 can be used for planar scans or single photon emission computed tomography (SPECT). Positron emitting 30 labels such as Fluorine-19 can be used in positron emission Paramagnetic ions such as Gadlinium (III) tomography. Manganese (II) can be used in magnetic resonance imaging (MRI). Localization of the label permits determination of the spread of the cancer. The amount of label within an organ or

- 15 -

tissue also allows determination of the presence or absence of cancer in that organ or tissue.

For patients diagnosed with prostate cancer, injection of an antibody which specifically binds CSG can also have a 5 therapeutic benefit. The antibody may exert its therapeutic effect alone. Alternatively, the antibody can be conjugated to a cytotoxic agent such as a drug, toxin or radionuclide to enhance its therapeutic effect. Drug monoclonal antibodies have been described in the art for example by Garnett and 10 Baldwin, Cancer Research 1986 46:2407-2412. The use of toxins conjugated to monoclonal antibodies for the therapy of various cancers has also been described by Pastan et al. Cell 1986 Yttrium-90 labeled monoclonal antibodies have 47:641-648. been described for maximization of dose delivered to the tumor 15 while limiting toxicity to normal tissues (Goodwin and Meares Other cytotoxic Cancer Supplement 1997 80:2675-2680). radionuclides including, but not limited to Copper-67, Iodine-131 and Rhenium-186 can also be used for labeling of antibodies against CSG.

20 Antibodies which can be used in these in vivo methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers single-stranded and oligonucleotides such as those derived from an in vitro 25 evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

Small molecules predicted via computer imaging to specifically bind to regions of CSGs can also be designed and synthesized and tested for use in the imaging and treatment of prostate cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to CSGs identified herein. Molecules identified in the library as being capable of binding to CSG are key candidates for further evaluation for use in the treatment of prostate cancer.

PCT/US99/24331

EXAMPLES

WO 00/23111

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments.

These exemplifications, while illustrating certain aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

All examples outlined here were carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following example can be carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Example 1: Identification of CSGs

Identification of CSGs were carried out by a systematic analysis of data in the LIFESEQ database available from Incyte Pharmaceuticals, Palo Alto, CA, using the data mining Cancer Leads Automatic Search Package (CLASP) developed by diaDexus LLC, Santa Clara, CA.

The CLASP performs the following steps: selection of highly expressed organ specific genes based on the abundance level of the corresponding EST in the targeted organ versus all the other organs; analysis of the expression level of each highly expressed organ specific genes in normal, tumor tissue, disease tissue and tissue libraries associated with tumor or disease; selection of the candidates demonstrating component ESTs were exclusively or more frequently found in tumor libraries. The CLASP allows the identification of highly expressed organ and cancer specific genes. A final manual in depth evaluation is then performed to finalize the CSGs selection.

- 17 -

Clones depicted in the following Table 1 are CSGs useful in diagnosing, monitoring, staging, imaging and treating prostate cancer.

Table 1: CSGs

5	Clone ID	Pro #	SEQ ID NO:
	3424528Н1	Pro109	1,2
	578349H1	Pro112	3,4
	1794013Н1	Proll1	5,6
	2189835H1	Prol15	7,8
10	3277219Н1	Prol10	9,10
	1857415	Prol13	11,12
	1810463H1	Proll4	13,14
	zr65G11	Proll8	15,16
	2626135H1		17
15	zd46d08		18
	1712252Н1		19
	784583H1		20

Example 2: Relative Quantitation of Gene Expression

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye.

25 During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control is used to 30 standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), ATPase, or 18S ribosomal RNA (rRNA) is used as this endogenous

control. To calculate relative quantitation between all the samples studied, the target RNA levels for one Sample were used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

The tissue distribution and the level of the target gene were evaluated for every sample in normal and cancer tissues. Total RNA was extracted from normal tissues, cancer tissues, and from cancers and the corresponding matched adjacent tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was done using primers and Taqman probes specific to each target gene. The results were analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the target gene in a particular tissue compared to the calibrator tissue.

Expression of Clone ID 3424528H1 (Pro109):

For the CSG Pro109, real-time quantitative PCR was 20 performed using the following primers:

Forward Primer:

5'- ATCAGAACAAAGAGGCTGTGTC - 3' (SEQ ID NO:21)
Reverse Primer:

5'- ATCTCTAAAGCCCCAACCTTC - 3' (SEQ ID NO:22)

25 The absolute numbers depicted in Table 2 are relative levels of expression of the CSG referred to as Pro109 in 12 normal different tissues. All the values are compared to normal stomach (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular 30 tissue from different individuals.

- 19 -

Table 2: Relative Levels of CSG Pro109 Expression in Pooled
Samples

Tissue	NORMAL
Colon	0.02
Endometrium	0.01
Kidney	0.48
Liver	14.83
Ovary	0.08
Pancreas	4.38
Prostate	11.24
Small Intestine	0.42
Spleen	0
Stomach	1
Testis	0.62
Uterus	0.02

5

10

15

The relative levels of expression in Table 2 show that with the exception of liver (14.83), Pro109 mRNA expression is higher (11.24) in prostate compared with all other normal tissues analyzed. Pancreas, with a relative expression level of 4.38, is the only other tissue expressing considerable mRNA for Pro109.

The absolute numbers in Table 2 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 3.

The absolute numbers depicted in Table 3 are relative levels of expression of Prol09 in 28 pairs of matching samples and 4 unmatched samples. All the values are compared to 30 normal stomach (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 3: Relative Levels of CSG Pro109 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro34B	Prostate 1	5.98	6.06
5	Pro65XB	Prostate 2	16.68	3.85
	Pro69XB	Prostate 3	20.46	6.82
	Pro78XB	Prostate 4	1.39	1.4
	Pro101XB	Prostate 5	24.8	9.8
	Pro12B	Prostate 6	9.1	0.2
10	Pro13XB	Prostate 7	0.5	9.7
	Pro20XB	Prostate 8	13	12.5
	Pro23B	Prostate 9	16.8	3
	Ovr100050	Ovary 1	0.4	
	Ovr1028	Ovary 2	1.9	
15	Ovr18GA	Ovary 3		0.1
	Ovr206I	Ovary 4		0.1
	Mam12X	Mammary Gland 1	13.5	1.4
	Mam47XP	Mammary Gland 2	0.7	0.2
	Lng47XQ	Lung 1	2.36	0.03
20	Lng60XL	Lung 2	7.39	0.2
	Lng75XC	Lung 3	0.77	0.27
	StoAC44	Stomach 1	0.05	1.19
	StoAC93	Stomach 2	0.55	0.8
	StoAC99	Stomach 3	0.12	3.04
25	ColAS43	Colon 1	16.11	0.07
	ColAS45	Colon 2	0.11	0.08
	ColAS46	Colon 3	4.99	0.4
	Liv15XA	Liver 1	8.43	10.97
	Liv42X	Liver 2	1.57	20.82

-	21	_
---	----	---

Liv94XA	Liver 3	2.98	9.19
Pan77X	Pancreas 1	36	32
Pan82XP	Pancreas 2	0.09	7.09
Pan92X	Pancreas 3	0.7	0
Pan71XL	Pancreas 4	2.48	0.73
Pan10343	Pancreas 5	46	5.5

0 = Negative

5

In the analysis of matching samples, the higher levels of expression were in prostate, showing a high degree of tissue specificity for prostate tissue. Of all the samples different than prostate analyzed, only 4 cancer samples (the cancer sample mammary 1 with 13.5, colon 1 with 16.11, liver 1 with 8.43, and lung 2 with 7.39) showed an expression comparable to the mRNA expression in prostate. These results confirmed some degree of tissue specificity as obtained with the panel of normal pooled samples (Table 2).

Furthermore, the level of mRNA expression was compared in cancer samples and the isogenic normal adjacent tissue from the same individual. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 3 shows overexpression of Prolog in 6 out of 9 primary prostate cancer tissues compared with their respective normal adjacents. Thus, overexpression in the cancer tissue was observed in 66.66% of the prostate matching samples tested (total of 9 prostate matching samples).

Altogether, the degree of tissue specificity, plus the mRNA overexpression in 66.66% of the primary prostate matching samples tested is indicative of Pro109 being a diagnostic 30 marker for prostate cancer.

- 22 -

Expression of Clone ID 578349H1 (Prol12):

For the CSG Proll2, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- TGCCGAAGAGGTTCAGTGC - 3' (SEQ ID NO:23)

Reverse Primer

5'- GCCACAGTGGTACTGTCCAGAT - 3' (SEQ ID NO:24)

The absolute numbers depicted in Table 4 are relative levels of expression of the CSG Proll2 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 4: Relative Levels of CSG Prol12 Expression in Pooled
Samples

Tissue	NORMAL
Brain	2.9
Heart	0.1
Kidney	0.2
Liver	0.2
Lung	7.7
Mammary	4.2
Muscle	0.1
Prostate	5.5
Small Intestine	1.8
Testis	1
Thymus	1
Uterus	21

20

15

5

25

The relative levels of expression in Table 4 show that 30 Proll2 mRNA expression is the 3rd most highly expressed gene (after uterus and mammary) in the pool of normal prostate tissue compared to a total of 12 tissues analyzed. The absolute numbers in Table 4 were obtained analyzing pools of samples of a particular tissue from different individuals.

35 These results demonstrate that Prol12 mRNA expression is specific for prostate thus indicating Prol12 to be a diagnostic marker for prostate disease especially cancer.

- 23 -

Expression of Clone ID 1794013H1 (Pro111):

For the CSG Proll1, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- GCTGCAAGTTCTCCACATTGA - 3' (SEQ ID NO:25)

Reverse Primer

5'- CAGCCGCAGGTGAAACAC - 3' (SEQ ID NO:26)

The absolute numbers depicted in Table 5 are relative levels of expression of the CSG Proll1 in 12 normal different tissues. All the values are compared to normal testis (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 5: Relative Levels of CSG Proll1 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.04
Heart	0
Kidney	0
Liver	0
Lung	0.05
Mammary	0.14
Muscle	5166.6
Prostate	1483.72
Small Intestine	0.33
Testis	1
Thymus	0.49
Uterus	0.07

25

20

15

5

The relative levels of expression in Table 5 show that Proll1 mRNA expression is extraordinarily high in the pool of normal prostate (1483.72) compared to all the other tissues analyzed with the exception of muscle (5166.6). These results demonstrate that Proll1 mRNA expression shows specificity for prostate and muscle.

The absolute numbers in Table 5 were obtained analyzing pools of samples of a particular tissue from different

- 24 -

individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples \mathfrak{F} f a single individual in Table 6.

The absolute numbers depicted in Table 6 are relative 5 levels of expression of Proll1 in 48 pairs of matching and 18 unmatched samples. All the values are compared to normal testis (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same 10 individual.

Table 6: Relative Levels of CSG Proll1 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro101XB	Prostate 1	8.3	21.8
15	Pro12B	Prostate 2	2336	133
	Pro13XB	Prostate 3	3.4	23
	Pro20XB	Prostate 4	21.6	121.5
	Pro23B	Prostate 5	19.4	3.7
	Pro34B	Prostate 6	15	39
20	Pro65XB	Prostate 7	8	867
	Pro69XB	Prostate 8	56	94
	Pro78XB	Prostate 9	24	1515
	Pro84XB	Prostate 10	119	15.35
	Pro90XB	Prostate 11	8.08	112.2
25	Pro91XB	Prostate 12	0.88	51.8
_	ProC215	Prostate 13	0.3	
	ProC234	Prostate 14	0.35	
	ProC280	Prostate 15	436.5	
	Pro109XB	Prostate 16	3.43	265
30	Proll0	Prostate 17	18.2	8.73

WO 00/23111

PCT/US99/24331

	Pro125XB	Prostate 18	0.34	186
	Pro326	Prostate 19	1392	110
	Pro10R	Prostate 20 (prostatitis)	0.5	
	Pro20R	Prostate 21 (prostatitis)	24.1	
5	Pro258	Prostate 22 (BPH)	4610	
	Pro263C	Prostate 23 (BPH)	0	
	Pro267A	Prostate 24 (BPH)	1.46	
	Pro271A	Prostate 25 (BPH)	0	
	Pro460Z	Prostate 26 (BPH)	1.47	
10	ProC032	Prostate 27 (BPH)	14.4	
	Tst39X	Testis 1	0	0
	Bld32XK	Bladder 1	0.44	0.41
	Bld46XK	Bladder 2	0	0
	Bld66X	Bladder 3	0	0
15	BldTR14	Bladder 4	0	0
	Kid106XD	Kidney 1	0	0
	Kid107XD	Kidney 2	0	0
	Kid109XD	Kidney 3	0	0
	Pan10343	Pancreas 1	0 .	0
20	Pan71XL	Pancreas 2	0	0
	Pan77X	Pancreas 3	0	0 -
	Liv15XA	Liver 1	0	0
	Liv42X	Liver 2	0	0
	ClnAS43	Colon 1	0	0 .
25	ClnAS45	Colon 2	0	0
	ClnAS46	Colon 3	0	0
	ClnAS67	Colon 4	0	0
	ClnAC19	Colon 5	0	0
	ClnAS12	Colon 6	0	0

- 26 -

	Intestine 2 0	0 0 0 0 .7 0
Q Lung 1	0	
		.7 0
L Lung 2		
		0 0
C Lung 3		0 0
Lung 4		0 0
Mammary	/ Gland 1	0 1.4
Mammary	/ Gland 2 0	.2 0
K Mammary	Gland 3	0 0
7 Mammary	Gland 4	0 0
<pre>Mammary</pre>	Gland 5	0
N Mammary	Gland 6	0
Covary 1	0.	14 0
Ovary 2	0.	. 2
Ovary 3)
OO Ovary 4	0.	. 2
Ovary 5		0
Ovary 6		0
Ovary 7		0.2
Ovary 8		0
	Lung 3 Lung 4 Mammary Mammary Mammary Mammary Mammary Mammary Movary 1 Ovary 2 Ovary 3 Ovary 4 Ovary 5 Ovary 6 Ovary 7	Lung 3 Lung 4 Mammary Gland 1 Mammary Gland 2 Mammary Gland 3 Mammary Gland 4 Mammary Gland 5 Mammary Gland 6 Movary 1 Ovary 2 Ovary 3 Ovary 3 Ovary 4 Ovary 5 Ovary 6 Ovary 7

0= Negative

In the analysis of matching samples, the higher levels of expression were in prostate showing a high degree of tissue specificity for prostate. These results confirm the tissue specificity results obtained with normal pooled samples (Table 5).

Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an 30 indication of specificity for cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 6 shows overexpression of Proll1 in 5 out

- 27 -

of 16 primary prostate cancer samples compared with their respective normal adjacent (prostate samples 2, 5, 10, 17, and 19). Similar expression levels were observed in 3 unmatched prostate cancers (prostate samples 13, 14, 15), 2 prostatitis (prostate samples 20, 21), and 6 benign prostatic hyperplasia samples (prostate samples 22 through 27). Thus, there is overexpression in the cancer tissue of 31.25% of the prostate matching samples tested (total of 16 prostate matching samples).

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 31.25% of the prostate matching samples tested are indicative of Proll1 being a diagnostic marker for prostate cancer.

Expression of Clone ID 2189835H1 (Pro115):

15 For the CSG Proll5, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- TGGCTTTGAACTCAGGGTCA - 3' (SEQ ID NO:27)

Reverse Primer

20 5'- CGGATGCACCTCGTAGACAG - 3' (SEQ ID NO:28)

The absolute numbers depicted in Table 7 are relative levels of expression of the CSG Prol15 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 7: Relative Levels of CSG Pro115 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.016
Heart	0.002
Kidney	8.08
Liver	2.20
Lung	112.99

30

_	20	_
_	~0	_

Mammary	29.45
Muscle	0.05
Prostate	337.79
Small Intestine	7.54
Testis	1.48
Thymus	1
Uterus	1.4

5

The relative levels of expression in Table 7 show that Prol15 mRNA expression is higher (337.79) in prostate compared 10 with all the other normal tissues analyzed. Lung, with a relative expression level of 112.99, and mammary (29.446) are the other tissues expressing moderate levels of mRNA for Prol15. These results establish Prol15 mRNA expression to be highly specific for prostate.

The absolute numbers in Table 7 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 8.

The absolute numbers depicted in Table 8 are relative levels of expression of Proll5 in 17 pairs of matching and 21 unmatched samples. All the values are compared to normal thymus (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the 25 normal adjacent sample for that same tissue from the same individual.

Table 8: Relative Levels of CSG Pro115 Expression in Individual Samples

-	Sample ID	Tissue	Cancer	Matching Normal Adjacent
30	Pro12B	Prostate 1	1475.9	190.3
	ProC234	Prostate 2	169.61	
	Pro109XB	Prostate 3		639.53
	Prol01XB	Prostate 4	1985.2	2882.9

- 29 -

WO 00/23111 PCT/US99/24331

	Pro13XB	Prostate 5	34.9	13.9
	Pro215	Prostate 6	525.59	
	Pro125XB	Prostate 7		556.05
	Pro23B	Prostate 8	1891.4	1118.6
5	ProC280	Prostate 9	454.3	
	Pro20XB	Prostate 10	1332.6	
	Pro34B	Prostate 11		362.91
	Pro65XB	Prostate 12		135.06
	Pro69XB	Prostate 13		179.67
10	Pro10R	Prostate 14 (prostatitis)	143.82	
	Pro20R	Prostate 15 (prostatitis)	397.79	
	Pro258	Prostate 16 (BPH)	216.6	
	Pro263C	Prostate 17 (BPH)	601.25	
	Pro267A	Prostate 18 (BPH)	200.28	
15	Pro271A	Prostate 19 (BPH)	111.43	
	Pro460Z	Prostate 20 (BPH)	53.84	
	ProC032	Prostate 21 (BPH)	56.94	
	SmI21XA	Small Intestine 1	28.8	29.9
	SmIH89	Small Intestine 2	70.8	348.5
20	ClnAC19	Colon 1	22.73	446.47
	ClnAS12	Colon 2	116.97	493.18
	Kid106XD	Kidney 1	86.13	41.14
	Kid107XD	Kidney 2	0.26	35.14
	Lng47XQ	Lung 1	5.13	20.98
25	Lng60XL	Lung 2	13.93	114.78
	Lng75XC	Lung 3	16.47	53.79
	Mam12X	Mammary Gland 1	6.25	10.75
	Mam162X	Mammary Gland 2	1.84	2.54
	Mam42DN	Mammary Gland 3	23.08	35.51

Ovr10050	Ovary 1	0.9	
Ovr1028	Ovary 2	261.4	o o
Ovr103X	Ovary 3	. 7	0.1
Ovr20GA	Ovary 4		0
Ovr25GA	Ovary 5		0

0 = Negative

5

Higher levels of expression were seen in prostate, showing a high degree of tissue specificity for prostate 10 tissue. Of all the analyzed samples different from prostate, only two cancer samples (colon 2 with 116.97 and ovary 2 with 261.4), and 5 normal adjacent tissue samples (small intestine 2, colon 1, colon 2, kidney 1, and lung 2), showed an expression comparable to the mRNA expression in prostate. 15 These results confirmed the tissue specificity results obtained with the panel of normal pooled samples (Table 7).

Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an 20 indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 8 shows higher expression of Proll5 in 3 out of 4 matched prostate cancer tissues (prostate samples 1, 5 & 8).

Altogether, the high level of tissue specificity, plus the higher expression in 75% of the prostate matching samples tested, are indicative of Proll5 being a diagnostic marker for prostate cancer.

Expression of Clone ID 3277219H1 (Prol10):

30 For the CSG ProllO, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- CGGCAACCTGGTAGTGAGTG - 3' (SEQ ID NO:29)

- 31 -

Reverse Primer

10

15

20

5'- CGCAGCTCCTTGTAAACTTCAG - 3' (SEQ I NO:30)

The absolute numbers depicted in Table 9 are relative levels of expression of the CSG ProllO in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 9: Relative Levels of CSG Prol10 Expression in Pooled Samples

Tissue	NORMAL
Brain	6.61
Heart	0.7
Kidney	0.74
Liver	7.94
Lung	11.88
Mammary	22.78
Muscle	6.77
Prostate	3.01
Small Intestine	1
Testis	2.58
Thymus	13.74
Uterus	2.61

The relative levels of expression in Table 9 show that Prol10 25 mRNA expression is not as high in normal prostate (3.01) compared with all the other normal tissues analyzed.

The absolute numbers in Table 9 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 10.

The absolute numbers depicted in Table 10 are relative levels of expression of Prol10 in 33 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from

- 32 -

the normal adjacent sample for that same tissue from the same individual. $\label{eq:sample}$

Table 10: Relative Levels of CSG Prol10 Expression in Individual Samples

5	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro12B	Prostate 1	11.8	0.3
	Pro78XB	Prostate 2	14.3	6.3
	Pro101XB	Prostate 3	33.2	10.7
	Pro13XB	Prostate 4	0.3	0.4
10	Pro23XB	Prostate 5	25.5	14.4
	Pro20XB	Prostate 6	43.3	4
	Pro34XB	Prostate 7	31.8	18.7
	Pro65XB	Prostate 8	26.9	3.4
	Pro69XB	Prostate 9	12.5	7
15	Lng75XC	Lung 1	1.9	3
	Lng90X	Lung 2	5.5	0.5
	LngAC11	Lung 3	9.3	9.7
	LngAC32	Lung 4	11.2	2.2
	Lng47XQ	Lung 5	11.3	0.3
20	Lng60XL	Lung 6	29.1	6.8
	Mam12B	Mammary Gland 1	19.8	0
	Mam603X	Mammary Gland 2	13.7	0
	Mam82XI	Mammary Gland 3	73.5	0
	MamA04	Mammary Gland 4	0	24.6
25	MamB011X	Mammary Gland 5	17.4	2
•	MamC012	Mammary Gland 6	0	12.8
	MamC034	Mammary Gland 7	0	61
Ī	Mam12X	Mammary Gland 8	14	2.2
	Mam59X	Mammary Gland 9	33	2.2

– 33 –

MamA06X	Mammary Gland 10	16.4	0.8
Liv15XA	Liver 1	4.7	0.6
Liv42X	Liver 2	7.5	2.6
Liv94XA	Liver 3	0.4	1.4
ClnAS43	Colon 1	52.9	1.4
ClnAS45	Colon 2	2.1	0.8
ClnAS46	Colon 3	39.8	3.7
SmI21X	Small Intestine 1	0.9	0.1
SmIH89	Small Intestine 2	5.8	0.9

 $10 \quad 0 = Negative$

5

The levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 10 shows overexpression of Prol10 in 8 of the 9 primary prostate cancer tissues compared with their respective normal adjacent (except prostate 4). Thus, there was overexpression in 88.88% of the cancer prostate tissue as compared to the prostate matching samples tested (total of 9 prostate matching samples).

Although not tissue specific, Prollo mRNA expression is upregulated in prostate cancer tissues. The mRNA overexpression in 88.88% of the primary prostate matching 25 cancer samples tested is indicative of Prollo being a diagnostic marker for prostate cancer. Prollo also showed overexpression in several other cancers tested including small intestine, colon, liver, mammary and lung (see Table 10). Accordingly Prollo may be a diagnostic marker for other types of cancer as well.

- 34 -

Expression of Clone ID 1857415; Gene ID 346880 (Pro113):

For the CSG Proll3, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- CGGGAACCTACCAGCCTATG - 3' (SEQ ID NO:31)

Reverse Primer

5'- CAGGCAACAGGGAGTCATGT - 3' (SEQ ID NO:32)

The absolute numbers depicted in Table 11 are relative levels of expression of the CSG Prol13 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 11: Relative Levels of CSG Prol13 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.03
Heart	0
Kidney	0.01
Liver	0
Lung	0
Mammary Gland	0
Muscle	0.04
Prostate	489.44
Small Intestine	0.02
Testis	0.35
Thymus	1
Uterus	0.13

The relative levels of expression in Table 11 show that Prol13 mRNA expression is higher (489.44) in prostate compared with all the other normal tissues analyzed. Testis, with a relative expression level of 0.35, uterus (0.13), thymus (1.0), kidney (0.01) and brain (0.03) were among the other tissues expressing lower mRNA levels for Prol13. These results establish that Prol13 mRNA expression is highly specific for prostate.

20

15

5

25

The absolute numbers in Table 11 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single 5 individual in Table 12.

The absolute numbers depicted in Table 12 are relative levels of expression of Proll3 in 78 pairs of matching and 25 unmatched tissue samples. All the values are compared to normal thymus (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In cancers (for example, ovary) where it was not possible to obtain normal adjacent samples from the same individual, samples from a different normal individual were analyzed.

Table 12: Relative Levels of CSG Prol13 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matched or Unmatched Normal Adjacent
	Pro780B/781B	Prostate 1	375.58	446.29
20	Pro1291B/1292B	Prostate 2	1060	31
	Pro139B96/140B96	Prostate 3	41	32
	Pro209B96/210B96	Prostate 4	505	255
	Pro1256B/1257B	Prostate 5	165.79	141.63
	Pro1293B/1294B	Prostate 6	1613.7	874.61
25	Pro694B/695B	Prostate 7	458.6	142.21
	Pro1012B/1013B	Prostate 8	1520	864
	Pro1222B/1223B	Prostate 9	939	530
	Pro845B/846B	Prostate 10	1552.4	374.6
	Pro1094B/1095B	Prostate 11	278.37	135.89
30	Pro650B/651B	Prostate 12	532.81	640.85

WO 00/23111

PCT/US99/24331

	Pro902B/903B	Prostate 13	609.05	415.86
	Pro916B/917B	Prostate 14	699.42	401.24
	Pro9821110A/110B	Prostate 15	156	487.8
	ProS9821326A/26B	Prostate 16	744.4	472.8
5	Pro9407c215	Prostate 17	1389.2	
	Pro9407c234	Prostate 18	305.5	
	Pro9407c280A	Prostate 19	894.5	
	Pro9409C010R	Prostate 20 (prostatitis)	269.7	
	Pro9404C120R	Prostate 21 (prostatitis)	299.2	
10	Pro1000258	Prostate 22 (BPH)	149.6	
	Pro4001263C	Prostate 23 (BPH)	576	
	Pro4001267A	Prostate 24 (BPH)	132.1	
	Pro9411C032	Prostate 25 (BPH)	118.2	
	Pro4001460Z	Prostate 26 (BPH)	276.3	-
15	Pro4001271A	Prostate 27 (BPH)	58.7	
	Kid1064D/65D	Kidney 1	0	0.1
	Kid1079D/1080D	Kidney 2	0.3	0.02
	Kid1097D/1098D	Kidney 3	35.14	0.32
	Kid1024D/1025D	Kidney 4	1.31	0
20	Kid1183D/1184D	Kidney 5	24.79	0
	Kid1242D/1243D	Kidney 6	0	0
-	Bld469K	Bladder 1		2.88
	Bld467K/468K	Bladder 2	2.65	
	Bld327K/328K	Bladder 3	0	4.05
25	Bld470K	Bladder 4		1.64
	Bld665T/664T	Bladder 5	0.21	1.99

	Bld1496K/1497K	Bladder 6	13.55	1.14
	Bld1721K/1722K	Bladder 7	120.16	1.34
	Tst239X/240X	Testis 1	31.5	0.73
	TstS9820647A/47B	Testis 2	15.7	0
5	TstS9820663A/663B	Testis 3	72	1.4
	SknS9821248A/248B	Skin 1	1.8	0.5
	SknS99448A/448B	Skin 2	251.6	0
	Skn99816A/816B	Skin 3	33	0.7
	Sto4004864A4/B4	Stomach 1	14.12	0
10	Sto4004509A3/B1	Stomach 2	40.74	39
	SmI9807A212A/213A	Small Intestine 1	0.1	0
	SmI9802H008/H009	Small Intestine 2	5.8	0.1
	Cln9608B012/B011	Colon 1	4.5	0
	Cln9709c074ra/073ra	Colon 2	65.8	3.1
15	Cln4004709A1/709B1	Colon 3	1.1	0.9
	Cln9405C199/C200	Colon 4	34.76	0.73
	Cln9707c004gb/006ga	Colon 5	90.26	0.96
	Cln96-09-B004/B003	Colon 6	17.9	20.64
	Cln9612B006/B005	Colon 7	17.56	0.3
20	Cln9705F002D/F001C	Colon 8	21.39	0
	ClnCXGA	Colon 9	429.14	142.69
	Pan10343a	Pancreas 1	0	0
	Pan776P/777P	Pancreas 2	0	0.15
	Pan9210/9220	Pancreas 3	7.36	0
25	Pan714L/715L	Pancreas 4	13.57	0.11
	Pan824P/825P	Pancreas 5	0	0
	Lng476Q/477Q	Lung 1	0	0
	Lng605L/606L	Lung 2	0	0.1
	Lng11145B/11145C	Lung 3	85.9	0

	Lng0008632A/32B	Lung 4	23.85	0
	Lng750C/751C	Lung 5	0.32	0.25
	Lng8890A/8890B	Lung 6	10.63	0
	Lng8926A/8926B	Lung 7	15.37	0
5	Lng0010239A/39B	Lung 8	26.17	0
	Lng9502C109R/110R	Lung 9	0.68	0
•	LngS9821944a/44b	Lung 10	0	0
	Mam00042D01/42N01	Mammary Gland 1	8.5	0
	Mam59XC	Mammary Gland 2	61.07	0 -
10	Mam9706A066G/67C	Mammary Gland 3	4.84	0
	Mam14153a1C	Mammary Gland 4	9.72	6.99
	Mam1620F/1621F	Mammary Gland 5	0.91	0 .
	Mam00014D05	Mammary Gland 6	2.45	0
	End10479B/D	Endometrium 1	133.43	1.12
15	End9705A125A/126A	Endometrium 2	0	0.39
	End9704C281A/282A	Endometrium 3	23.5	1.56
	End680097/681097	Endometrium 4	88.89	79.02
	Utr13590/13580	Uterus 1	0.2	0
	Utr850U/851U	Uterus 2	0	0
20	Utr14170/14180	Uterus 3	14	0.4
	Utr233U96/234U96	Uterus 4	8.65	4.64
	CvxVNM00052D01/52N01	Cervix 1	0.82	77.15
	CvxVNM00083D01/83N01	Cervix 2	0.78	221.48
	CvxND00023D01/23N01	Cervix 3	3.25	15.22
25	Ovr10370/10380	Ovary 1	0.1	0
	Ovr10050	Ovary 2	18.96	
	Ovr1028	Ovary 3	0	
	Ovr14638A1C	Ovary 4	3.2	
	Ovr14603A1D	Ovary 5	882.3	
30	0vr7730	Ovary 6	0	

Ovr9702C018GA	Ovary 7	0.15
Ovr206I	Ovary 8	0
Ovr9702C020GA	Ovary 9	0
Ovr9702C025GA	Ovary 10	0
Ovr9701C035GA	Ovary 11	0.07
Ovr9701C050GB	Ovary 12	0.58

0 = Negative

5

In the analysis of matching samples, the higher levels of expression were in prostate, showing a high degree of 10 tissue specificity for prostate tissue. In addition to the higher expression levels in prostate cancer samples, Prol13 expression was found to be either induced (where not expressed in normal adjacent tissues) or somewhat upregulated in several other cancers. However, the relative expression and the fold increase in prostate cancer samples far exceeds that in other cancer tissues and is highly significant.

Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an 20 indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 12 shows overexpression of Prol13 in 13 out of 16 primary prostate cancer tissues compared with their respective normal adjacent (prostate samples 2, 3, 4, 5, 6 7, 25 8, 9, 10, 11, 13, 14, 16). Thus, there was overexpression in the cancer tissue for 81.25% of the prostate matching samples The median for the level of expression in prostate cancer tissue samples is 609, whereas the median for all other cancers is only 7.93, with the exception of one colon sample, 30 colon 9, whose expression was similar to that found in prostate cancer tissues.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 81.25% of the primary prostate matching samples tested are indicative of Proll3 being a

- 40 -

diagnostic marker for prostate cancer. Expression was also found to be higher in other cancer tissues compared with their respective normal adjacent tissues (kidney, bladder, testis, skin, stomach, small intestine, colon, pancreas, lung, mammary, endometrium, uterus, and ovary) thus indicating Prol13 to be a pan cancer marker.

Expression of Clone ID 1810463H1 (Pro114):

For the CSG Proll4, real-time quantitative PCR was performed using the following primers:

10 Forward Primer

5'- TGGGCATCTGGGTGTCAA - 3' (SEQ ID NO:33)

Reverse Primer

5'- CGGCTGCGATGAGGAAGTA - 3' (SEQ ID NO:34)

The absolute numbers depicted in Table 13 are relative levels of expression of the CSG Proll4 in 12 normal different tissues. All the values are compared to normal muscle (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

20 Table 13: Relative Levels of CSG Proll4 Expression in Pooled Samples

	Tissue	NORMAL
	Brain	9.7
	Heart	0.7
;	Kidney	414.4
	Liver	4
	Lung	882.2
	Mammary	44
	Muscle	1
)	Prostate	1951
	Small Intestine	22
	Testis	367.1
	Thymus	25.8
	Uterus	139.6

35 The relative levels of expression in Table 13 show that Proll4 mRNA expression is higher (1951) in prostate compared with all the other normal tissues analyzed. Lung, with a relative

25

30

- 41 -

expression level of 882.2, kidney 414.4, testis 367.1 and uterus 139.6, are the other tissues expressing himsher levels of mRNA for Prol14. These results establish Prol14 mRNA expression to be more specific for prostate than other tissues 5 examined.

The high level of tissue specificity is indicative of Proll4 being a diagnostic marker for diseases of the prostate, especially cancer.

Expression of Clone ID zr65g11 (Prol18):

10 For the CSG Prol18, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- GCCCATCTCCTGCTTCTTTAGT - 3' (SEQ ID NO:35)

Reverse Primer

15

5' - CGTGGAGATGGCTCTGATGTA - 3' (SEQ ID NO:36)

The absolute numbers depicted in Table 14 are relative levels of expression of the CSG Prol18 in 12 normal different tissues. All the values are compared to normal kidney (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 14: Relative Levels of CSG Pro118 Expression in Pooled Samples

	Tissue	NORMAL	
25	Colon	0.87	
	Endometrium	19282	1
	Kidney	1	1
	Liver	0	1
	Ovary	86.22	
30	Pancreas	0	
	Prostate	962.1	1
	Small Intestine	0	
	Spleen	0.75	1
	Stomach	0.54	1
35	Testis	343.7	1
	Uterus	1064	1

- 42 -

The relative levels of expression in Table \$4 show that Prol18 mRNA expression is the 3rd highest in prostate (962.1) next to endometrium (19282) and uterus (1064), which are female-specific tissues. Testis, with a relative expression level of 343.7 is the only other male tissue expressing moderate levels of mRNA for Prol18. These results establish Prol18 mRNA expression to be highly specific for reproductive tissues including the prostate.

The absolute numbers in Table 14 were obtained analyzing 10 pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 15.

The absolute numbers depicted in Table 15 are relative 15 levels of expression of Proll8 in 59 pairs of matching and 21 unmatched samples. All the values are compared to normal kidney (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same 20 individual.

Table 15: Relative Levels of CSG Prol18 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro12B	Prostate 1	41700.7	22242.83
5	ProC234	Prostate 2	40087	
	Pro78XB	Prostate 3	4075.6	7066.7
	Pro109XB	Prostate 4	334.4	777.2
	Pro84XB	Prostate 5	11684	58290
	Pro101XB	Prostate 6	21474.13	100720.8
0	Pro91X	Prostate 7	14849	33717
	Pro13XB	Prostate 8	202.57	146.91

25

30

	ProC215	Prostate 9	73243	
	Pro125XB	Prostate 10	629.6	521.4
	Pro23B	Prostate 11	157532.6	110654.4
	Pro90XB	Prostate 12	2317	64134
5	ProC280	Prostate 13	42020	
	Pro20XB	Prostate 14	2909.31	
	Pro34B	Prostate 15	29610	23264
	Pro110	Prostate 16	13354	30991
	Pro65XB	Prostate 17	10126	11270
10	Pro69XB	Prostate 18		2671.42
	Pro326	Prostate 19	9962.3	19231
	Pro10R	Prostate 20 (prostatitis)	27355	
	Pro20R	Prostate 21 (prostatitis)	21081	
	Pro258	Prostate 22 (BPH)	79916.32	
15	Pro263C	Prostate 23 (BPH)	108924.5	
	Pro267A	Prostate 24 (BPH)	92910.22	
	Pro271A	Prostate 25 (BPH)	57004.4	
	Pro460Z	Prostate 26 (BPH)	57449.23	
	ProC032	Prostate 27 (BPH)	45781.44	
20	Kid106XD	Kidney 1	3.08	217.36
	Kid107XD	Kidney 2	0	38.36
	Kid109XD	Kidney 3	0	123.5
	Kid10XD	Kidney 4	17.69	67.8
	Kid11XD	Kidney 5	16.74	360.8
25	Kid124D	Kidney 6	0	167.4
	Bld32XK	Bladder 1	0	0
	Bld47K	Bladder 2		36.38
	Bld66X	Bladder 3	0	4.52
	BldTR14	Bladder 4	0	12.17

	BldTR17	Bladder 5	0	0
	Bld46XK	Bladder 6	16.5	0
	Tst39X	Testis 1	116.6	24.35
	Tst647T	Testis 2	856.16	43.5
5	StoAC44	Stomach 1	0	0
	StoAC93	Stomach 2	0	0
	SmI21XA	Small Intestine 1	68.45	0
	SmIH89	Small Intestine 2	0	0
	ClnAC19	Colon 1	149	21.33
10	ClnAS12	Colon 2	0	0
	ClnB34	Colon 3	0	0
	ClnB56	Colon 4	13.04	5.22
	ClnAS43	Colon 5	0	0
	Lng47XQ	Lung 1	0	0 .
15	Lng60XL	Lung 2	0	0
	Lng75XC	Lung 3	0	3.38
	Lng90X	Lung 4	0	0
	LngBR26	Lung 5	0	26.82
	Pan10343	Pancreas 1	50.47	0
20	Pan77X	Pancreas 2	281.1	0
	Pan92X	Pancreas 3	18.41	0
	Pan71XL	Pancreas 4	0	0
	Pan82XP	Pancreas 5	0	0
	PanC044	Pancreas 6	0	0
25	Mam12X	Mammary Gland 1	0	0
	Mam162X	Mammary Gland 2	0	0
•	Mam42DN	Mammary Gland 3	0	0
	MamS127	Mammary Gland 4	12.58	0
	Mam14DN	Mammary Gland 5	0	0
30	End28XA	Endometrium 1	331.9	1824

	End3AX	Endometrium 2	27825	65839
	End4XA	Endometrium 3	10.3	15935
	Utr1410	Uterus 1	18885	18116
	Utr23XU	Uterus 2	3358	7674
5	CvxKS52	Cervix 1	0	0
	CvxKS83	Cervix 2	0	0
	Ovr10050	Ovary 1	72.86	
	Ovr1028	Ovary 2	0	
	Ovr638A	Ovary 3	0	
10	Ovr63A	Ovary 4	90.88	
	Ovr7730	Ovary 5	1.21	
	Ovr10400	Ovary 6	5.08	
	Ovr1050	Ovary 7	0	
	Ovr1118	Ovary 8	7.41	:
15	Ovr103X	Ovary 9		32.78
	Ovr20GA	Ovary 10		0 .
	Ovr25GA	Ovary 11		1173.83
	Ovr35GA	Ovary 12		313.4
	Ovr50GB	Ovary 13		823.1
20	Ovr18GA	Ovary 14		40.6
	Ovr206I	Ovary 15		1264
	Ovr230A	Ovary 16		1285

0 = Negative

In the analysis of matching samples, the higher levels of expression were in prostate, endometrium, testis, and ovary showing a high degree of tissue specificity for reproductive tissues. These results confirmed the tissue specificity results obtained with the panel of normal pooled samples (Table 14).

30 Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an

- 46 -

indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 15 shows overexpression of Prol18 in 5 out of 14 primary prostate cancer tissues (prostate samples 1, 8, 10, 11, 15) compared with their respective normal adjacent. Thus, there was overexpression in the cancer tissue for 35.71% of the prostate matching samples tested (total of 14 prostate matching samples). Expression of Prol18 was similarly higher in 3 unmatched cancer tissues (prostate samples 9, 13, 14), 2 prostatitis (prostate samples 20, 21), and 6 benign hyperplasia tissues (prostate samples 22 through 27).

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 35.71% of the primary prostate matching samples tested are indicative of Proll8 being a diagnostic marker for prostate cancer.

- 47 -

What is claimed is:

1. A method for diagnosing the presence of prostate cancer in a patient comprising:

- (a) determining levels of CSG in cells, tissues or bodily 5 fluids in a patient; and
- (b) comparing the determined levels of CSG with levels of CSG in cells, tissues or bodily fluids from a normal human control, wherein a change in determined levels of CSG in said patient versus normal human control is associated with the 10 presence of prostate cancer.
 - 2. A method of diagnosing metastases of prostate cancer in a patient comprising:
 - (a) identifying a patient having prostate cancer that is not known to have metastasized;
- (b) determining CSG levels in a sample of cells, tissues, or bodily fluid from said patient; and
- (c) comparing the determined CSG levels with levels of CSG in cells, tissue, or bodily fluid of a normal human control, wherein an increase in determined CSG levels in the 20 patient versus the normal human control is associated with a cancer which has metastasized.
 - 3. A method of staging prostate cancer in a patient having prostate cancer comprising:
 - (a) identifying a patient having prostate cancer;
- 25 (b) determining CSG levels in a sample of cells, tissue, or bodily fluid from said patient; and
- (c) comparing determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in determined CSG levels in said patient versus the normal human control is associated with a cancer which is progressing and a decrease in the determined CSG levels is associated with a cancer which is regressing or in remission.

- 4. A method of monitoring prostate cancer in a patient for the onset of metastasis comprising:
- (a) identifying a patient having prostate cancer that is not known to have metastasized;
- (b) periodically determining levels of CSG in samples of cells, tissues, or bodily fluid from said patient; and
- (c) comparing the periodically determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the 10 periodically determined CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.
 - 5. A method of monitoring a change in stage of prostate cancer in a patient comprising:
- 15 (a) identifying a patient having prostate cancer;
 - (b) periodically determining levels of CSG in cells, tissues, or bodily fluid from said patient; and
- (c) comparing the periodically determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the periodically determined CSG levels in the patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease is associated with a cancer which is regressing in stage or in remission.
- 6. A method of identifying potential therapeutic agents for use in imaging and treating prostate cancer comprising screening molecules for an ability to bind to CSG wherein the ability of a molecule to bind to CSG is indicative of the molecule being useful in imaging and treating prostate cancer.
- 7. The method of claim 1, 2, 3, 4, 5 or 6 wherein the CSG comprises SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12,

- 49 -

- 13, 14, 15, 16, 17, 18, 19 or 20 or a polypeptide encoded thereby.
 - 8. An antibody which specifically binds CSG.
- A method of imaging prostate cancer in a patient
 comprising administering to the patient an antibody of claim
 8.
 - 10. The method of claim 9 wherein said antibody is labeled with paramagnetic ions or a radioisotope.
- 11. A method of treating prostate cancer in a patient 10 comprising administering to the patient an antibody of claim 7.
 - 12. The method of claim 11 wherein the antibody is conjugated to a cytotoxic agent.

SEQUENCE LISTING

```
<110> Salceda, Susana
       Recipon, Herve
       Cafferkey, Robert
       diaDexus, LLC
 <120> Method of Diagnosing, Monitoring, Staging, Imaging and
       Treating Prostate Cancer
 <130> DEX-0052
 <140>
 <141>
 <150> 60/104,737
<151> 1998-10-19
<160> 36
<170> PatentIn Ver. 2.0
<210> 1
<211> 188
<212> DNA
<213> Homo sapiens
<400> 1
ggtaaacacc tgcttttatc atcagaacaa agaggctgtg tcccctgccc tatgaggtcc 60
atttctgaga gttgtggcta atgggcaaga aggttggggc tttagagatt tgggataaag 120
atatcaaaca ccagaaaggt agaaagaagt gatcagatta gggttactta ggtgatgata 180
tgaactct
                                                                   188
<210> 2
<211> 9819
<212> DNA
<213> Homo sapiens
<400> 2
cagctggggt ctacccaggt ccatgtcttg gacatgttga gagtttttct ggaaggcagg 60
gatacagtgt ggtccaaaaa cacacaaatg cccctactgg cccaggggtt gtcacaatag 120
actggaaggg tgacacatcc caggcgcttg ccacccatca cacgcacctc ctacccactg 180
gcatccttcc accccaggca cacacaaagc ctcagtccag agatcaactc tggactcagc 240
totgaatttg catatootgt gtgtagattc attottcata acctotgccc agcotagott 300
gtgtatcatt ttttttctc tattagggga ggagcccgtc ctggcactcc cattggcctg 360
tagattcacc tecectggge agggeeccag gaceeaggat aatatetgtg cetectgeec 420
agaaccetce aageagacae aatggtaaga atggtgeetg teetgetgte tetgetgetg 480
cttctgggtc ctgctgtccc ccaggagaac caagatggtg agtggggaaa gcaagggatg 540
```

ggtgctggag	g aggactgga	a ggaggtgagg	g aacaggacat	gtggctggg	a gacaggetge	600
atgcagctg	g gataccctg	g catacggcag	g gaatgggtgo	ccaaggctgt	caactccctc	660
agctcacaca	a cttccagga	g cattcaggga	gcctctgcgc	tggcccgaaa	taagacctto	720
aggaatctga	a atctaaaac	cctagtttac	: agtgaaaaca	aagactccaa	agaccaagcg	780
acctgcttgg	g ggtagacagi	caggacggag	, taggaaccat	atgcctggag	, ctgcttctgc	840
		c cgatggctgg				
		g tgggaaggca				
		a ctcagacaco				
		tccgggagat				
		gtgcaggata				
		catttacaaa				
		, ttaatctgat				
		taggtggtac				
		tggatacaca				
		gagatttatt				
		agagccggcc				
		ggcagactgt				
		gccttcaact				
		tactgatttt				
		tggggtacat				
		tgaggtatcc				
agacaatcca	attatacttt	tttggttatt	ttagttttta	aaagtatttg	attatttatt	1860
tatttattta	tttttgagac	agagtctcac	tctgtcaccc	aggcaggagt	gcagtggcat	1920
		tccgcctccc				
ctgagtagct	aggactacag	gcacctgcca	ccacacctgg	ctaattttt	tgtattttta	2040
gtagagacgg	tttcatcatg	ttggccaggc	tagtcttgat	atcctgacct	cgtgatctgc	2100
ccgccttggt	ctcccaaagt	gccgggatta	caggtgtcag	caactgcgcc	tggcctctct	2160
tttggttatt	taaaagtgta	caattaaatt	atgattatta	ttattattt	tgagatggat	2220
tcttgttctg	tcacccaggc	tggagtgcag	tggcgtgatc	ttggcttact	gcaaacctcc	2280
gcctgttggg	ttcaagcaat	tatcttgcct	cgggtgtaca	ctgccacaca	cggctaactt	2340
atgtatttt	aatagagata	gggtttcacc	atgttggcta	gactggtctt	gacctcttga	2400
cctcaagtga	tccactcact	tcagcctccc	agagtgctgg	aattacaggc	acgagecace	2460
acacctggcc	ccagttaaat	tattattgac	tatagtcacc	ctgttgtgct	atcaaatagt	2520
aggtcttatt	cattcttctt	tttttttt	tttttgtgac	agagttgccc	aggctggaat	2580
gcagtggtgc	aatcttggct	cactgcaacc	tctgcctccc	gggcttaagc	gattctcctg	2640
cctcagcctt	ctgagtcgct	gggactacag	gtgtgtgcca	ccacgcccgg	ctaatttatg	2700
tatttttagt	agagatgggg	tttcaccatg	ttggccaggc	tggtttcgaa	ctcctgacct	2760
caagtgaccc	acctgcctca	gcttcccaaa	gtgttggaat	tacaggcatg	agccaccaca	2820
cctggcccca	gttaaattat	tattcactgg	agtcactttg	ttgtgctatc	aaatagtttt	2880
ctaactattt	tttttgtacc	cattaaccac	cctcccaatt	tccccccaac	cctgccacta	2940
cccttcccag	cctttggtaa	ccatccttct	actctctatg	tccatgaatt	caattgtagg	3000
gtctactgat	ttaaaggcta	atcacattta	gacactcagg	agcaagaata	attttagtaa	3060
ttgaactagg	attctgccat	atgacctcca	acatcattag	cacctgtgta	aattgtatca	3120
taaaataatt	atggaactat	tatggaaatg	tccctctctc	ccagatccca	ccttgtacca	3180
aaatgcaagg	tacaaccccg	ggaattctga	gctccatcct	agtcttaccc	tgtgctaatt	3240
cagtctgggt	catttcttga	attttctggt	aaattctcct	ttctaccctt	tctaactata	3300
tgtatttgtc	aggttaagct	agaagtgtta	atttttttt	tttttgagat	ggagccttgc	3360
tttgtcacct	aggctgaagt	gcagtggcat	gatctcagct	cactgcaagc	teegeeteee	3420

gggttcatg	c catteteet	g cctcagcct	c ctgagtagc	t gggactaca	g gcacccgcc	a 3480
	g ctaattttt					
	a tctcctgac					
	a gccactgag					
	a tccaagctgg					
	a agtgatttt					
	c cagctaattt					
aggctggtc	t tcaactcctg	g atctcaagta	a atctgcctg	c cttggcctc	caaagtcctg	3900
ggattacag	g catgagccad	ggagcccag	c ctagaaatg	t taatttctaa	a cgcatgtcag	3960
attccatgo	a cactgggcaa	ggttccatto	ctccatggg	g tgactcaggg	g atccaggcca	a 4020
	t gagactcttt					
	a aagggaaagc					
	t ctcttttgac					
	t tatcttccta					
	a atatcatctc					
	a gttgatggca					
	cctatgcagc					
	gcaggtcgtt					
	cccgcgtttc					
	a gacaggaagt					
	g caggacagcc					
	gtggagtatt					
	aacccaagag					
	caatcccagg					
	ccccacagca aggtcacaca					
	tccacacccc					
	ccagggaggt					
	gtggttctgc					
	tgcaaacatt					
	gttctatgaa					
	gggtacagcc					
	acacaccgta					
	gagttccttg.					
	ctgcgctatg					
ctgtgccccc	ggcccagtgc	tcttctcct	aaaccacact	gccagcccca	aggcagccaa	5580
cctcaggtct	ggtgaactgc	tggtgttaaa	ttatcataga	gtgggtgtca	aaagatgggc	5640
tactaagtac	aaaaatgccc	aaggtgctac	atgggatctg	aagattttca	aaaggaggca	5700
	aggcagatgt					
	gtgtgtcagc					
	ttgatggaca					
	gaacacaggg					
	agcactttgg					
	gccaacatgg					
	acgtctgtgg					
	cagaggttac					
	cgttgtctaa					
tgacatttac	tgagcactta	ctttgtgcca	ggcccatcta	tgagcatata	taatgctcag	6300

aatagcccc	taaaacagtg	ctcttggcat	tgccatttca	a gaggtgagga	a aatagaggca	6360
	g agtggctcca					
	a cgtgagccca					
accagtgct	g ggtaaacacc	tgcttttato	atcagaacaa	a agaggctgtg	tecectgee	6540
	atttctgaga					
	g atatcaaaca					
	a tgaactcttc					
	a ataatttcta					
cctcaggact	: cagcataaat	tcaaacaaat	aattgtcctt	agaagagatg	, ctatagaaga	6840
gatagaaata	tattcatatt	ctgtagcttt	ttttttttg	, agatggagtt	ttgctcttgt	6900
cacccaagct	ggagtgcagt	gatgcaatct	cagctcactg	caaactttgc	ctcctgggtt	6960
caagggatto	tcctgcctca	gcctcccgat	aactgggact	acaggetaca	ggcatgtgtc	7020
actactcct	gttaattttt	tttttttt	tttaagactg	agtettgete	tgtctttcag	7080
gctgatgtac	: aatggctcca	tctcggctca	ctacaacttc	tgtcccccag	gttcaagcga	7140
ttctcctgcc	tcagcctcat	gagtagctgg	gattacaggo	atgtgccago	acacccagca	7200
aatttttgta	tttttagtag	agatgaggtc	ttaccatgtt	ggccaggctg	gtctcaaact	7260
cctgacctca	ggtgatcctt	tggcctcagc	ctccctaact	gctgggatta	caggcatgag	7320
ccactgcgtc	cagcctaatt	ttatatttt	ggtagagatg	gggtttcacc	atattggcca	7380
ggctggtctc	gaactcatga	cctaaggtga	tccatcctcc	tcagcctctc	aaagtgctgg	7440
gattacaagt	gtgagccact	gggcctggtg	cttttttt	tttttttt	tttttttt	7500
tgagataggg	tctcactctg	tcacccaggc	tgaaatgcag	tagtgtgatt	ttggctcatt	7560
gcagccttga	cttcccaggc	tgaagtgatc	ctcccacctc	agcctcctga	gtagctgggg	7620
ctacaggcat	gcaccaccat	gctgcgctaa	tttttatatt	ttttgtagtg	gtgggatttc	7680
gccatatcac	cctggctggt	ctggaacccc	tgggctcaag	cgatccactc	gcttcagctt	7740
ctcaaagtgc	tgggattaca	ggcatgagcc	acagcgccca	ggctgtagct	ctcttaagga	7800
ggaacatatc	tcatctgaga	caaacctgaa	atgccaaacc	aaactgagtt	agcccctctc	7860
tgtctgttgt	atatattgga	gtaataacct	atttgtcttg	ataaagggat	tgcatgcttg	7920
aattgcaaaa	acctttattt	cttttgggtt	gcccaatgtg	caagactaag	agttattttg	7980
ataaatttct	caccaggctg	actgtctctc	tgtggggtcg	ggggagtttt	cagggtctca	8040
cgtattgcag	ggaaggtttg	gttgtgagat	cgagaataac	agaagcagcg	gagcattctg	8100
gaaatattac	tatgatggaa	aggactacat	tgaattcaac	aaagaaatcc	cagcctgggt	8160
ccccttcgac	ccagcagccc	agataaccaa	gcagaagtgg	gaggcagaac	cagtctacgt	8220
gcagcgggcc	aaggcttacc	tggaggagga	gtgccctgcg	actctgcgga	aatacctgaa	8280
atacagcaaa	aatatcctgg	accggcaagg	tactcactgc	ttcctgctcc	ccagtactga	8340
gcccagaata	aaagacgatc	tcaggctagg	agctcaggca	acatcttagt	ccggtctcat	8400
ctgttcctgg	atgtccctca	gacccccagc	tttcatcttt	taggatttat	tccttccctg	8460
	atttgtggtc					
	ctttcttgat					
	gaggtgaggg					
	ttccaaatcc					
ggtggagaga	agggaggcgt	gtaaaagtca	gccccagaag	gacaagagca	agccagtgtg	8760
	atggctgcaa					
	cagggaagtg					
	ggacagaact					
	acaacttatt					
	cgtcagcctt					
	cttcgagctc					
ctgggccggg	agaagtccac	ctcaagcctg	cagtgtcaca	ctctatccct	ccacagatcc	9180

```
tccctctgtg gtggtcacca gccaccaggc cccaggagaa aagaagaaac tgaagtgcct 9240
ggcctacgac ttctacccag ggaaaattga tgtgcactgg actcgggccg gcgaggtgca 9300
  ggagcctgag ttacggggag atgttcttca caatggaaat ggcacttacc agtcctgggt 9360
ggtggtggca gtgcccccgc aggacacagc cccctactcc tgccacgtgc agcacagcag 9420
  cctggcccag ccctcgtgg tgccctggga ggccagctag gaagcaaggg ttggaggcaa 9480
  tgtgggatet cagacccagt agetgeeett cetgeetgat gtgggagetg aaccacagaa 9540
  atcacagtca atggatccac aaggcctgag gagcagtgtg gggggacaga caggaggtgg 9600
  atttggagac cgaagactgg gatgcctgtc ttgagtagac ttggacccaa aaaatcatct 9660
caccttgage ccacceceae eccattgtet aatetgtaga agetaataaa taateateee 9720
  teettgeeta geataacaga gaateetttt tttaaeggtg atgegetgta gaaatgtgae 9780
  tagattttct cattggttct gccctcaagc actgaattc
 <210> 3
  <211> 250
  <212> DNA
  <213> Homo sapiens
 <400> 3
 cgccctgcg ccgccgagcc agctgccaga atgccgaact ggggaggagg caagaaatgt 60
 ggggtgtgtc agaagacggt ttactttgcc gaagaggttc agtgcgaagg caacagcttc 120
 cataaatcct gcttcctgtg catggtctgc aagaagaatc tggacagtac cactgtggcc 180
 gtgcatggtg aggagattta ctgcaagtcc tgctacggca agaagtatgg gcccaaaggc 240
 tatggctacg
 <210> 4
 <211> 1900
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> unsure
 <222> (16)
 <220>
 <221> unsure
 <222> (18)
 <220>
 <221> unsure
 <222> (20)
 <220>
 <221> unsure
 <222> (1887)
 <220>
 <221> unsure
```

<222> (1894)

```
<400> 4
 acgcettecg eggagnanan caaaacggeg egeaggeegg gegeacceag eegecaette 60
 egagagegee tgeegeeet ggegeegeeg ageeagetge cagaatgeeg aactggggag 120
 gaggcaagaa atgtggggtg tgtcaagaag acggtttact ttgccgaaga ggttcagtgc 180
 gaaggcaaca gcttccataa atcctgcttc ctgtgcatgg tctgcaagaa gaatctggac 240
 agtaccactg tgggccgtgc atggtgagga gatttactgg caagtccctg ctacggcaag 300
 aagtatgggc ccaaaggcta tggctacggg ccagggcgca ggcaccctca gcactgacaa 360
 gggggagteg etgggtatea ageaegagga ageeeetggg eeaeaggeee accaecaace 420
 ccaatggcat ccaaatttgc ccagaagatt ggtggctccg agcgctgccc ccgatgcagc 480
 caggeagtet atgetgegga gaaggtgatt ggtgetggga agteetggea taaggeetge 540
 tttcgatgtg ccaagtgtgg caaaggcctt gagtcaacca ccctgggcag acaaggatgg 600
 cgagatttac tgcaaaggat gttatgctaa aaacttcggg cccaagggct ttggttttgg 660
 gcaaggaget ggggeettgg tecaetetga gtgaggeeae cateaeceae cacaecetge 720
ccactcctgc gcttttcatc gccattccat tcccagcagc tttggagacc tccaggatta 780
tttetetgte agecetgeea catateaeta atgaettgaa ettgggeate tggeteeett 840
tggtttgggg gtctgcctga ggtcccaccc cactaaaggg ctccccaggc ctgggatctg 900
acaccatcac cagtaggaga cotcagtgtt ttgggtctag gtgagagcag gcccctctcc 960
ccacacctcg ccccacagag ctctgttctt agcctcctgt gctgcgtgtc catcatcage 1020
tgaccaagac acctgaggac acatcttggc acccagagga gcagcagcaa caggctggag 1080
ggagagggaa gcaagaccaa gatgaggagg ggggaaggct gggttttttg gatctcagag 1140
atteteetet gtgggaaaga ggttgagett eetggtgtee eteagagtaa geetgaggag 1200
tcccagctta gggagttcac tattggaggc agagaggcat gcaggcaggg tcctaggagc 1260
ccctgcttct ccaggcctct tgcctttgag tctttgtgga atggatagcc tcccactagg 1320
actgggagga gaataaccca ggtcttaagg accccaaagt caggatgttg tttgatcttc 1380
tcaaacatet agtteeetge ttgatgggag gateetaatg aaatacetga aacatatatt 1440
ggcatttatc aatggctcaa atcttcattt atctctggcc ttaaccctgg ctcctgaggc 1500
tgcggccagc agagcccagg ccagggctct gttcttgcca cacctgcttg atcctcagat 1560
gtggagggag gtaggcactg cctcagtctt catccaaaca cctttccctt tgccctgaga 1620
cctcagaatc ttccctttaa cccaagaccc tgcctcttcc actccaccct tctccaggga 1680
cccttagatc acatcactcc acccctgcca ggccccaggt taggaatagt ggtgggagga 1740
aggggaaagg gctgggcctc accgctccca gcaactgaaa ggacaacact atctggagcc 1800
acceactgaa agggetgeag geatgggetg tacceaaget gattteteat etggteaata 1860
                                                                   1900
aagctgttta gaccagaaaa aaaaaanaaa aaanaaaagg
<210> 5
<211> 273
<212> DNA
<213> Homo sapiens
<400> 5
gatgcatcaa aagagctgca agttctccac attgacttct tgaatcagga caacgccgtt 60
teteaceaca catgggagtt ccaaacgage agteetgtgt teeggegagg acaggtgttt 120
cacctgcggc tggtgctgaa ccagccccta caatcctacc accaactgaa actggaattc 180
agcacaggge cgaatcctag catcgccaaa cacaccctgg tggtgctcga cccgaggacg 240
ccctcagacc actacaactg gcaggcaacc ctt
                                                                  273
<210> 6
```

<211> 3021 <212> DNA <213> Homo sapiens

<400> 6 tgtggaagca ccaggcatca gagatagagt cttccctggc attgcaggag agaatctgaa 60 gggatgatgg atgcatcaaa agagctgcaa gttctccaca ttgacttctt gaatcaggac 120 aacgccgttt ctcaccacac atgggagttc caaacgagca gtcctgtgtt ccggcgagga 180 caggigittic accigegget ggigetgaac cagecectae aatectaeca ccaacigaaa 240 ctggaattca gcacagggcc gaatcctagc atcgccaaac acaccctggt ggtgctcgac 300 ccgaggacgc cctcagacca ctacaactgg caggcaaccc ttcaaaatga gtctggcaaa 360 gaggtcacag tggctgtcac cagttccccc aatgccatcc tgggcaagta ccaactaaac 420 gtgaaaactg gaaaccacat ccttaagtct gaagaaaaca tcctatacct tctcttcaac 480 ccatggtgta aagaggacat ggttttcatg cctgatgagg acgagcgcaa agagtacatc 540 ctcaatgaca cgggctgcca ttacgtgggg gctgccagaa gtatcaaatg caaaccctgg 600 aactttggtc agtttgagaa aaatgtcctg gactgctgca tttccctgct gactgagagc 660 teceteaage ceacagatag gagggaeece gtgetggtgt geagggeeat gtgtgetatg 720 atgagetttg agaaaggeea gggegtgete attgggaatt ggaetgggga etatgaaggt 780 ggcacagece catacaagtg gacaggcagt geeegatee tgcagcagta etacaacaeg 840 aagcaggctg tgtgctttgg ccagtgctgg gtgtttgctg ggatcctgac tacagtgctg 900 agagegttgg geateceage aegeagtgtg acaggetteg atteagetea egacacagaa 960 aggaacctca cggtggacac ctatgtgaat gagaatggca agaaaatcac cagtatgacc 1020 cacgactotg totggaattt coatgtgtgg acggatgcct ggatgaagcg accggatctg 1080 cccaagggct acgacggctg gcaggctgtg gacgcaacgc cgcaggagcg aagccagggt 1140 gtcttctgct gtgggccatc accactgacc gccatccgca aaggtgacat ctttattgtc 1200 tatgacacca gattcgtctt ctcagaagtg aatggtgaca ggctcatctg gttggtgaag 1260 atggtgaatg ggcaggagga gttacacgta atttcaatgg agaccacaag catcgggaaa 1320 aacatcagca ccaaggcagt gggccaagac aggcggagag atatcaccta tgagtacaag 1380 tatccagaag gctcctctga ggagaggcag gttcatggat catgccttcc tccttctcag 1440 ttctgagagg gagcacagac gacctgtaaa agagaacttt cttcacatgt cggtacaatc 1500 agatgatgtg ctgctgggaa actctgttaa tttcaccgtg attcttaaaa ggaagaccgc 1560 tgccctacag aatgtcaaca tcttgggctc ctttgaacta cagttgtaca ctggcaagaa 1620 gatggcaaaa ctgtgtgacc tcaataagac ctcgcagatc caaggtcaag tatcagaagt 1680 gactotgacc ttggactoca agacctacat caacagootg gotatattag atgatgagoo 1740 agttatcaga ggtttcatca ttgcggaaat tgtggagtct aaggaaatca tggcctctga 1800 agtattcacg tetttecagt accetgagtt etetatagag ttgeetaaca caggeagaat 1860 tggccagcta cttgtctgca attgtatctt caagaatacc ctggccatcc ccttgactga 1920 egtcaagtte tetttggaaa geetgggeat etecteaeta cagaeetetg accatgggtg 1980 agtotgcctg aggacggtgc agcctggtga gaccatccaa tcccaaataa aatgcacccc 2040 aatgctcaga agattgttct catcaccaag tagccttgtc tgatgctgtg gagccttagt 2160 tgagatttca gcatttccta ccttgtggct tagctttcag attatggatg attaaatttg 2220 atgacttata tgagggcaga ttcaagagcc agcaggtcaa aaaggccaac acaaccataa 2280 gcagccagac ccacaaggcc aggtcctgtg ctatcacagg gtcaccttct tttacagtta 2340 gaaacaccag ccgaggccac agaatcccat ccctttcctg agtcatggcc tcaaaaatca 2400 999ccaccat tgtctcaatt caaatccata gatttcgaag ccacagattc tctccctgga 2460 gcaagcatga ctatgggcag cccagtgctg ccacctgctg acgacccttg agaagctgcc 2520 atatetteag gecatgggtt caccageeet gaaggeaeet gteaaetgga gtgetetete 2580

```
agcactggga tgggcctgat agaagtgcat teteeteeta ttgeeteeat teteetetet 2640
 ctatccctga aatccaggaa gtccctctcc tggtgctcca agcagtttga agcccaatct 2700
 gcaaggacat ttctcaaggg ccatgtggtt ttgcagacaa ccctgtcctc aggcctgaac 2760
 tcaccataga gacccatgtc agcaaacggt gaccagcaaa tcctcttccc ttattctaaa 2820
 getgeeeett gggagaetee agggagaagg cattgettee teeetggtgt gaactettte 2880
 tttggtattc catccactat cctggcaact caaggctgct tctgttaact gaagcctgct 2940
 cettettgtt etgeceteca gagatttget caaatgatea ataagettta aattaaacte 3000
 tacttcaaga aaaaaaaacc g
                                                                   3021
 <210> 7
 <211> 267
 <212> DNA
 <213> Homo sapiens
<400> 7
gaacattcca gatacctatc attactcgat gctgttgata acagcaagat ggctttgaac 60
tragggtrae carragetat tggacettae tatgaaaace atggatacca accggaaaac 120
ccctatcccg cacagcccac tgtggtcccc actgtctacg aggtgcatcc ggctcagtac 180
taccegtece cegtgeecca gtacgeeceg agggteetga egeaggette caacceegte 240
gtctgcacgc agcccaaatc cccatcc
                                                                   267
<210> 8
<211> 3443
<212> DNA
<213> Homo sapiens
<400> 8
gggcgggccg ggccgagtag gcgcgagcta agcaggaggc ggaggcggag gcggagggcg 60
aggggegggg agegeeget ggagegege aggteatatt gaacatteca gatacetate 120
attactegat getgttgata acagcaagat ggetttgaac teagggteac caccagetat 180
tggaccttac tatgaaaacc atggatacca accggaaaac ccctatcccg cacagcccac 240
tgtggtcccc actgtctacg aggtgcatcc ggctcagtac tacccgtccc ccgtgcccca 300
gtacgccccg agggtcctga cgcaggcttc caaccccgtc gtctgcacgc agcccaaatc 360
cccatccggg acagtgtgca cctcaaagac taagaaagca ctgtgcatca ccttgaccct 420
ggggaccttc ctcgtgggag ctgcgctggc cgctggccta ctctggaagt tcatgggcag 480
caagtgctcc aactctggga tagagtgcga ctcctcaggt acctgcatca acccctctaa 540
ctggtgtgat ggcgtgtcac actgccccgg cggggaggac gagaatcggt gtgttcgcct 600
ctacggacca aacttcatcc ttcaggtgta ctcatctcag aggaagtcct ggcaccctgt 660
gtgccaagac gactggaacg agaactacgg gcgggcggcc tgcagggaca tgggctataa 720
gaataatttt tactctagcc aaggaatagt ggatgacagc ggatccacca gctttatgaa 780
actgaacaca agtgccggca atgtcgatat ctataaaaaa ctgtaccaca gtgatgcctg 840
ttcttcaaaa gcagtggttt ctttacgctg tatagcctgc ggggtcaact tgaactcaag 900
ccgccagagc aggatcgtgg gcggcgagag cgcgctcccg ggggcctggc cctgggcagg 960
teagestgea egtecagaac gtecaegtgt geggaggete cateateace ceegagtgga 1020
tcgtgacagc cgcccactgc gtggaaaaac ctcttaacaa tccatggcat tggacggcat 1080
ttgcggggat tttgagacaa tctttcatgt tctatggagc cggataccaa gtagaaaaag 1140
tgatttotca tocaaattat gactocaaga coaagaacaa tgacattgog otgatgaago 1200
tgcagaagcc tctgactttc aacgacctag tgaaaccagt gtgtctgccc aacccaggca 1260
```

```
tgatgctgca gccagaacag ctctgctgga tttccgggtg gggggccacc gaggagaaag 1320
  ggaagacete agaagtgetg aacgetgeea aggtgettet cattgagaca cagagatgea 1380
 acagcagata tgtctatgac aacctgatca caccagccat gatctgtgcc ggcttcctgc 1440
 aggggaacgt cgattcttgc cagggtgaca gtggagggcc tctggtcact tcgaagaaca 1500
 atatctggtg gctgataggg gatacaagct ggggttctgg ctgtgccaaa gcttacagac 1560
 caggagtgta cgggaatgtg atggtattca cggactggat ttatcgacaa atgagggcag 1620
 acggctaatc cacatggtct tegteettga egtegtttta caagaaaaca atggggetgg 1680
 ttttgcttcc ccgtgcatga tttactctta gagatgattc agaggtcact tcatttttat 1740
 taaacagtga acttgtctgg ctttggcact ctctgccatt ctgtgcaggc tgcagtggct 1800
 cccctgccca gcctgctctc cctaacccct tgtccgcaag gggtgatggc cggctggttg 1860
 tgggcactgg cggtcaagtg tggaggagag gggtggaggc tgccccattg agatcttcct 1920
 gctgagtcct ttccaggggc caattttgga tgagcatgga gctgtcacct ctcagctgct 1980
 ggatgacttg agatgaaaaa ggagagacat ggaaagggag acagccaggt ggcacctgca 2040
 geggetgeet etggggeeae ttggtagtgt ecceageeta ectetecaea aggggatttt 2100
 gctgatgggt tcttagagcc ttagcagccc tggatggtgg ccagaaataa agggaccagc 2160
 ccttcatggg tggtgacgtg gtagtcacct tgtaagggga acagaaacat ttttgttctt 2220
 atggggtgag aatatagaca gtgcccttgg gtgcgaggga agcaattgaa aaggaacttg 2280
 ccctgagcac tcctggtgca ggtctccacc tgcacattgg gtggggctcc tgggagggag 2340
 acteageett ceteeteate effectgace effeteetag caccetggag agtgeacatg 2400
 ccccttggtc ctgggcaggg gcgccaagtc tggcaccatg-ttggcctctt caggcctgct 2460
 agtcactgga aattgaggtc catgggggaa atcaaggatg ctcagtttaa ggtacactgt 2520
 ttccatgtta tgtttctaca cattgctacc tcagtgctcc tggaaactta gcttttgatg 2580
 totocaagta gtocacctto atttaactot tigaaactgt atcatotttg ccaagtaaga 2640
 gtggtggcct atttcagctg ctttgacaaa atgactggct cctgacttaa cgttctataa 2700
 atgaatgtgc tgaagcaaag tgcccatggt ggcggcgaag aagagaaaga tgtgttttgt 2760
 tttggactet etgtggteee ttecaatget gtgggtttee aaccagggga agggteeett 2820
 ttgcattgcc aagtgccata accatgagca ctactctacc atggttctgc ctcctggcca 2880
agcaggctgg tttgcaagaa tgaaatgaat gattctacag ctaggactta accttgaaat 2940
ggaaagtett gcaateeeat ttgcaggate egtetgtgea catgeetetg tagagageag 3000
cattcccagg gaccttggaa acagttggca ctgtaaggtg cttgctcccc aagacacatc 3060
ctaaaaggtg ttgtaatggt gaaaacgtct tccttcttta ttgccccttc ttatttatgt 3120
gaacaactgt ttgtcttttt ttgtatcttt tttaaactgt aaagttcaat tgtgaaaatg 3180
aatatcatgc aaataaatta tgcgattttt ttttcaaagt aaccactgca tctttgaagt 3240
tetgeetggt gagtaggace ageeteeatt teettataag ggggtgatgt tgaggetget 3300
99tcagagga ccaaaggtga ggcaaggcca gacttggtgc tcctgtggtt ggtgccctca 3360
gtteetgeag cetgteetgt tggagaggte ceteaaatga eteetteta ttattetatt 3420
agtctgtttc catgggcgtg ata
                                                                   3443
<210> 9
<211> 254
<212> DNA
<213> Homo sapiens
<400> 9
gtgctgcacc aggccaccat cetgcccaag actgggacag tgtccctgga ggtacggctc 60
ctggaggcct cccgtgcctt cgaggtgtca gagaacggca acctggtagt gagtgggaag 120
gtgtaccagt gggatgaccc tgaccccagg ctcttcgacc acccggaaag ccccaccccc 180
aaccccacgg agcccctctt cctggcccag gctgaagttt acaaggagct gcgtctgcgt 240
```

```
ggctacgact acgg
                                                                    254
 <210> 10
 <211> 8470
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> unsure
 <222> (4131)
 <220>
 <221> unsure
 <222> (5117)
<220>
<221> unsure
 <222> (5552)
<400> 10
eggeegtega caeggeageg geeeeggeet eeeteteege egegetteag eeteeegete 60
egeegegete cageeteget eteegeegee egeaeegeeg ceegegeeet caceagagea 120
gccatggagg aggtggtgat tgccggcatg tccgggaagc tgccagagtc ggagaacttg 180
caggagttet gggacaacet categgeggt gtggacatgg teaeggacga tgacegtege 240
tggaaggcgg ggctctacgg cctgcccgg cggtccggca agctgaagga cctgtctagg 300
tttgatgcct ccttcttcgg agtccacccc aagcaggcac acacgatgga ccctcagctg 360
eggetgetge tggaagteac etatgaagee ategtggaeg gaggeateaa eceagattea 420
ctccgaggaa cacacactgg cgtctgggtg ggcgtgagcg gctctgagac ctcggaggcc 480
ctgagccgag accccgagac actcgtgggc tacagcatgg tgggctgcca gcgagcgatg 540
atggccaacc ggctctcctt cttcttcgac ttcagagggc ccagcatcgc actggacaca 600
geetgeteet ceageetgat ggeeetgeag aacgeetace aggeeateea cagegggeag 660
tgccctgccg ccatcgtggg gggcatcaat gtcctgctga agcccaacac ctccgtgcag 720
ttettgagge tggggatget cageceegag ggeacetgea aggeettega caeagegggg 780
aatgggtact gccgctcgga gggtgtggtg gccgtcctgc tgaccaagaa gtccctggcc 840
cggcgggtgt acgccaccat cctgaacgcc ggcaccaata cagatggctt caaggagcaa 900
ggcgtgacct teceeteagg ggatateeag gageagetea teegetegtt gtaceagteg 960
gccggagtgg cccctgagtc atttgaatac atcgaagccc acggcacagg caccaaggtg 1020
ggcgaccccc aggagctgaa tggcatcacc cgagccctgt gcgccacccg ccaggagccg 1080
etgeteateg getecaceaa gtecaacatg gggeaceegg agecageete ggggetggea 1140
gecetggeea aggtgetget gteeetggag eaegggetet gggeeecaa eetgeaette 1200
catagececa accetgagat eccagegetg ttggatggge ggetgeaggt ggtggaceag 1260
eccetgeeg teegtggegg caacgtggge atcaacteet ttggettegg gggeteeaaa 1320
egtgeacate atcetgagge ccaacaegea geegeeeeee geaceeggee cacatgeeae 1380
cctgccccgt ctgctgcggg ccagcggacg cacccctgag gccgtgcaga agctgctgga 1440
gcagggcctc cggcacagcc agggcctggc tttcctgagc atgtgaacga catcgcggct 1500
gtccccgacc accgccatgc ccttccgtgg ctacgctgtg ctgggtggtg agacgcggtg 1560
geocagaggt geageaggtg ecegetggeg agegeeeget etggtteate tgetetggga 1620
tgggcacaca gtggcgcggg atggggctga gcctcatgcg cctggaccgc ttccgagatt 1680
```

ccatcctac	g ctccgatgag	gctgtgaaco	gattcggcct	gaaggtgtca	a cagctgctgc	1740
tgagcacag	a cgagagcacc	tttgatgaca	a tcgtccatto	gtttgtgag	c ctgactgcca	1800
tccagatag	g cctcatagac	ctgctgagct	gcatggggct	gaggccagat	ggcatcgtcg	1860
gccactccc	t gggggaggtg	gcctgtggct	acgccgacgg	ctgcctgtcd	: caggaggagg	1920
ccgtcctcg	tgcctactgg	aggggacagt	gcatcaaaga	agcccatcto	ccgccgggcg	1980
	c cgtgggcttg					
	c cacaactcca					
	g gagcagctga					
	cactcctact					
	c cgggagccga					
	g cacagcagcc					
ggtgagccct	gtgctgttcc	aggaggccct	gtggcacgtg	cctgagcacg	cggtggtgct	2400
	g ccccacgccc					
	ccctgatga					
	ctgcacctct					
	gctccccgag					
cctggcctgg	gacgcgccgg	ccgccgagga	cttccccaac	ggttcaggtt	cccctcagc	2700
caccatctac	acatgcacac	caagctccga	gtctcctgac	cgctacctgg	tggaccacac	2760
catcgacggt	cgcgtcctct	tccccgccac	tggctacctg	agcatagtgt	ggaagacgct	2820
ggcccgaccc	ctgggcctgg	gcgtcgagca	gctgcctgtg	gtgtttgagg	atgtggtgct	2880
gcaccaggco	accatcctgc	ccaagactgg	gacagtgtcc	ctggaggtac	ggctcctgga	2940
ggcctcccgt	gccttcgagg	tgtcagagaa	cggcaacctg	gtagtgagtg	ggaaggtgta	3000
ccagtgggat	gaccctgacc	ccaggctctt	cgaccacccg	gaaagcccca	ccccaaccc	3060
cacggagccc	ctcttcctgg	cccaggctga	agtttacaag	gagctgcgtc	tgcgtggcta	3120
cgactacggc	cctcatttcc	agggcatcct	ggaggccagc	ctggaaggtg	actcggggag	3180
gctgctgtgg	aaggataatg	ggtgagttca	tggacaccat	gctgcagatg	tccatcctgg	3240
gtcggccaag	cacggcctgt	acctgcccac	ccgtgtcacc	gccatccaca	tegaccetge	3300
cacccacagg	cagaagctgt	acacactgca	ggacaaggcc	caagtggctg	acgtggtggt	3360
gagcaggtgg	ctgagggtca	cagtggccgg	aggcgtccac	atctccgggc	tccacactga	3420
gtcggccccg	cggcggcagc	aggagcagca	ggtgcccatc	ctggagaagt	tttgcttcac	3480
	gaggaggggt					
gtgcaagggg	ctggtcgagg	cactcgagac	caaggtgacc	cagcaggggc	tgaagatggt	3600
	ctggatgggg					
	ggctgcctgc					
	ccaggagagg					
	actcaaggcc					
	ggaggtgctg					
	tcccctgctg					
	ccaggccgag					
	ccccagcgcc					
	ggacccgcct					
	ctgcacacac					
	gagccgcagt					
	gtgtcgctgc					
	cgccggccca					
	tgggtggagt					
	gaaggccatc					
tccgccgaga	gcccggcgga	acgctccggt	gtgtgctgct	ctccaacctc	agcagcacct	4560

cccacgtcc	c ggaggtgga	c ccgggctcc	g cagaactgc	a gaaggtgtt	g cagggagac	c 4620	
tggtgatga	a cgtctaccg	c gacggggcct	gggggctt	t ccgccactt	c ctgctggag	g 4680	
aggacaagc	c tgaggagcc	g acggcacate	g cctttgtga	g caccctcac	c cggggggac	c 4740	
tgtccctcc	a tccgctggg	t ctgctcctcg	g ctgcgccate	g cccagccca	c ctgccctgg	c 4800	
gcccagctc	t gcacggtct	a ctacgcctco	ctcaacttc	c gcgacatca	gctggccac	t 4860	
ggcaagctg	t cccctgatg	c catcccaggg	g aagtggacc	t cccaggaca	g cctgctagg	t 4920	
atggagttc	t cgggccgaga	a cgccagcgg	aagcgtgtg	a tgggactgg	gcctgccaa	g 4980	
ggcctggcc	a cctctgtcc	getgtcaceg	gacttcctc	t gggatgtgc	ttccaactg	g 5040	
acgctggag	g aggcggcct	ggtgcctgtd	gtctacage	a cggcctacta	a cgcgctggt	g 5100	
gtgcgtggg	gggtgeneed	cggggagacg	ctgctcatco	actcgggct	gggcggcgtg	g 5160	
ggccaggccg	g ccatcgccat	cgccctcagt	ctgggctgcd	gcgtcttcad	caccgtggg	g 5220	
tcggctgaga	a agcgggcgta	cctccaggcc	aggttcccc	agetegacag	caccagette	5280	
gccaactcc	gggacacato	cttcgagcag	catgtgctgt	ggcacacggg	cgggaaggg	5340	
gttgacctgg	g tettgaacte	: cttggcggaa	gagaagctgo	aggccagcgt	gaggtgcttg	5400	
gctacgcacg	g gtcgcttcct	ggaaattggc	aaattcgacc	tttctcagaa	ccacccgcto	5460	
ggcatggcta	tcttcctgaa	gaacgtgaca	ttccacgggg	, tcctactgga	tgcgttcttc	5520	
aacgagagca	gtgctgactg	gcgggaggtg	tnggcgcttg	tgcaggccgg	catccgggat	5580	
ggggtggtad	ggcccctcaa	gtgcacggtg	ttccatgggg	cccaggtgga	ggacgcctto	5640	
cgctacatgg	, cccaagggaa	. gcacattggc	aaagtcgtcg	tgcaggtgct	tgcggaggag	5700	
ccggaggcag	tggctgaagg	gggccaaacc	caagctgatg	teggecatet	ccaagacctt	5760	
ctgcccggcc	: cacaagagct	acatcatcgc	tggtggtctg	ggtggcttcg	gcctggagtt	5820	
ggcgcagtgg	ctgatacago	gtggggtgca	gaagctcgtg	ttgacttctc	gctccgggat	5880	
ccggacaggc	taccaggcca	agcaggtccg	ccggtggagg	cgccagggcg	tacaggtgca	5940	
		gctcactgga					
		ggcgtcttca					
		ttcttccagg					
		cgagggcgtg					
		gcaatgcggg					
		aaacgccggc					
		ggcattttgg					
		cgcatggcgt					
		agcagctttg					
		ctggtggagg					
		agctcactgg					
		gagcgtgagc					
		ctgcaggagc					
		ggaggatggt					
		ggagggcccc					
		tggtgcaccc					
		atccccacct					
		gctgcctact					
		ggctactcct					
		agcccagccc					
		gcctacaccc					
		gaggccatat					
		gcgctgctgc					
		aagagccacc					
ggcccggtc	cttctactac	aagctgcgtg	ccgctgagca	gtacacaccc	aaggccaagt	7440	

```
accatggcaa cgtgatgcta ctgcgcgcca agacgggtgg cgcctacggc gaggacctgg 7500
 gcgcggacta caacctctcc caggtatgcg acgggaaagt atccgtccac gtcatcgagg 7560
 gtgaccaccg cacgetgetg gagggeageg geetggagte cateateage ateatecaca 7620
 getecetgge tgagecaege gtgagegtge gggagggeta ggeeegtgee eeegeetgee 7680
 accggaggte actecaccat coccaccca teccaccca coccegecat gcaacgggat 7740
 tgaagggtcc tgccggtggg accetgtccg gcccagtgcc actgccccc gaggctagct 7800
 agacgtaggt gttaggcatg tcccacccac ccgccgcctc ccacggcacc tcggggacac 7860
 cagagetgee gaettggaga etectggtet gtgaagagee ggtggtgeee gtgeeegeag 7920
 gaactggggc tgggcctcgt gcgcccgtgg ggtctgcgct tggtctttct gtgcttggat 7980
 ttgcatattt attgcattgc tggtagagac ccccaggcct gtccaccctg ccaagactcc 8040
 traggrageg tgtgggtrer gractriger recattirer egatgirere tgrgggregeg 8100.
 ggcagccacc caagcetget ggetgeggee eeetetegge caggeattgg eteageeege 8160
 tgagtggggg gtcgtgggcc agtccccgag gactgggccc ctgcacaggc acacagggcc 8220
 eggccacace cageggeece eegcacagee accegtgggg tgetgeeett atgeeeggeg 8280
 ccgggcacca actccatgtt tggtgtttgt ctgtgtttgt ttttcaagaa atgattcaaa 8340
 ttgctgcttg gattttgaaa tttactgtaa ctgtcagtgt acacgtctgg accccgtttc 8400
 atttttacac caatttggta aaaatgctgc tctcagcctc ccacaattaa accgcatgtg 8460
 atctccaaaa
 <210> 11
<211> 812
<212> DNA
<213> Homo sapiens
<400> 11
gccgcagcca atcagcgcgc gtgcccgggc ccctgcgtct cttgcgtcaa gacggccgtg 60
ctgagcgaat gcaggcgact tgcgagctgg gagcgattta aaacgctttg gattcccccg 120
gcctgggtgg ggagagcgag ctgggtgccc cctagattcc ccgccccgc acctcatgag 180
ccgaccctcg gctccatgga gcccggcaat tatgccacct tggatggagc caaggatatc 240
gaaggettge tgggageggg aggggggegg aatetggteg cecaetecee tetgaceage 300
cacccagegg egectaeget gatgeetget gteaactatg ecceettgga tetgeeagge 360
teggeggage gecaaageaa tgecacecat geeetggggt geeecagggg aegteeccag 420
etecegtgee ttatggttae tttggaggeg ggtactaete etgeegagtg teeeggaget 480
egetgaaacc etgtgeeeag geageeacce tggeegegta eecegeggag acteeeacgg 540
ccggggaaga gtaccccagc cgccccactg agtttgcctt ctatccggga tatccgggaa 600
cctaccagec tatggccagt tacctggacg tgtctgtggt gcagactctg ggtgctcctg 660
gagaaccgcg acatgactcc ctgttgcctg tggacagtta ccagtcttgg gctctcgctg 720
gtggctggaa cagccagatg tgttgccagg gagaacagaa cccaccaggt cccttttgga 780
aggcagcatt tgcagactcc agcgggcagc ac
                                                                  812
<210> 12
<211> 2385
<212> DNA
<213> Homo sapiens
<400> 12
ataagetggg gtaaagtatt ttegeagttt etgeetttag gattttatta gettetetee 60
cccaggccgc agccaatcag cgcgcgtgcc cgggcccctg cgtctcttgc gtcaagacgg 120
```

```
ccgtgctgag cgaatgcagg cgacttgcga gctgggagcg atttaaaacg ctttggattc 180
 ccccggcctg ggtggggaga gcgagctggg tgccccctag attccccgcc cccgcacctc 240
 atgageegae ceteggetee atggageeg geaattatge cacettggat ggageeaagg 300
 atatcgaagg ettgetggga gegggagggg ggeggaatet ggtegeeeae teeeetetga 360
 ccagccaccc agcggcgcct acgctgatgc ctgctgtcaa ctatgccccc ttggatctgc 420
 caggetegge ggageegeea aageaatgee acceatgeee tggggtgeee caggggaegt 480
 ecccagetee egtgeettat ggttaetttg gaggegggta etaeteetge egagtgteee 540
 ggageteget gaaaccetgt geceaggeag ceaecetgge egegtaeeee geggagaete 600
 ccacggccgg ggaagagtac cccagccgcc ccactgagtt tgccttctat ccgggatatc 660
 cgggaaccta ccagcctatg gccagttacc tggacgtgtc tgtggtgcag actctgggtg 720
 ctcctggaga accgcgacat gactccctgt tgcctgtgga cagttaccag tcttgggctc 780
 tegetggtgg etggaacage cagatgtgtt gecagggaga acagaaceca ecaggteeet 840
 tttggaaggc agcatttgca gactccagcg ggcagcaccc tcctgacgcc tgcgcctttc 900
 gtcgcggccg caagaaacgc attccgtaca gcaaggggca gttgcgggag ctggagcggg 960
 agtatgegge taacaagtte atcaccaagg acaagaggeg caagateteg geagecacea 1020
 gcctctcgga gcgccagatt accatctggt ttcagaaccg ccgggtcaaa gagaagaagg 1080
 ttctcgccaa ggtgaagaac agcgctaccc cttaagagat ctccttgcct gggtgggagg 1140
agcgaaagtg ggggtgtcct ggggagacca ggaacctgcc aagcccaggc tggggccaag 1200
gactotgotg agaggoocot agagacaaca coottoocag gocactggot gotggactgt 1260
tcctcaggag cggcctgggt acccagtatg tgcagggaga cggaacccca tgtgacagcc 1320
cactccacca gggttcccaa agaacctggc ccagtcataa tcattcatcc tgacagtggc 1380
aataatcacg ataaccagta ctagctgcca tgatcgttag cctcatattt tctatctaga 1440
gctctgtaga gcactttaga aaccgctttc atgaattgag ctaattatga ataaatttgg 1500
aaggcgatcc ctttgcaggg aagctttctc tcagaccccc ttccattaca cctctcaccc 1560
tggtaacagc aggaagactg aggagagggg aacgggcaga ttcgttgtgt ggctgtgatg 1620
teegtttage attitietea getgacaget gggtaggtgg acaattgtag aggetgtete 1680
ttcctccctc cttgtccacc ccatagggtg tacccactgg tcttggaagc acccatcctt 1740
aatacgatga tttttctgtc gtgtgaaaat gaagccagca ggctgcccct agtcagtcct 1800
tccttccaga gaaaaagaga tttgagaaag tgcctgggta attcaccatt aatttcctcc 1860
cccaaactet etgagtette cettaatatt tetggtggtt etgaccaaag caggteatgg 1920
tttgttgagc atttgggatc ccagtgaagt agatgtttgt agccttgcat acttagccct 1980
teccaggeae aaacggagtg geagagtggt gecaaccetg tttteccagt ceaegtagae 2040
agattcacgt gcggaattct ggaagctgga gacagacggg ctctttgcag agccgggact 2100
ctgagaggga catgagggcc tetgcctetg tgttcattet etgatgteet gtacetggge 2160
teagtgeeeg gtgggaetea teteetggee gegeageaaa geeagegggt tegtgetggt 2220
cottectgca cottaggetg ggggtggggg gcctgccggc gcattetcca cgattgagcg 2280
cacaggeetg aagtetggae aaceegeaga acegaagete egageagegg gteggtggeg 2340
agtagtgggg tcggtggcga gcagttggtg gtgggccgcg gccgc
<210> 13
<211> 221
<212> DNA
<213> Homo sapiens
<400> 13
dsdnrstate tttetgtgtg gtgeageeet gttggeagtg ggeatetggg tgteaatega 60
tggggcatcc tttctgaaga tcttcgggcc actgtcgtcc agtgccatgc agtttgtcaa 120
```

egtgggetae tteeteateg cageeggegt tgtggtettt getettggtt teetgggetg 180

```
ctatggtgct aagactgaga gcaagtgtgc cctcgtgacg t
                                                                  221
 <210> 14
 <211> 1533
 <212> DNA
 <213> Homo sapiens
 <400> 14
 gggcacgcag acattetggg aagccacttg coccaccct gggctgcttc ttettgagat 60
 caggaggggc gttgcccagg gctggtgttg ccaggtggag gcctgctgag gcagtggttg 120
 tggggatcgg tetecaggea geaggggea geagggteaa ggagaggeta aetggeeaeg 180
 ggtggggcca gcaggcgggc agaaggaggc tttaaagcgc ctaccctgcc tgcaggtgag 240
 cagtggtgtg tgagagccag gccgtccctc tgcctgccca ctcagtggca acacccggga 300
 gctgttttgt cctttgtgga gcctcagcag ttccctgctt tcagaactca ctgccaagag 360
 ccctgaacag gagccaccat ggcagtgctt cagcttcatt aagaccatga tgatcctctt 420
caatttgctc atctttctgt gtggtgcagc cctgttggca gtgggcatct gggtgtcaat 480
cgatggggca tcctttctga agatcttcgg gccactgtcg tccagtgcca tgcagtttgt 540
caacgtgggc tacttcctca tcgcagccgg cgttgtggtc tttgctcttg gtttcctggg:600
etgetatggt getaagaetg agageaagtg tgeectegtg aegttettet teateeteet 660
ceteatette attgetgagg ttgcagetge tgtggtegee ttggtgtaca ccacaatgge 720
tgagcacttc ctgacgttgc tggtagtgcc tgccatcaag aaagattatg gttcccagga 780
agacttcact caagtgtgga acaccaccat gaaagggctc aagtgctgtg gcttcaccaa 840
ctatacggat tttgaggact caccctactt caaagagaac agtgcctttc ccccattctg 900
ttgcaatgac aacgtcacca acacagccaa tgaaacctgc accaagcaaa aggctcacga 960
ccaaaaagta gagggttgct tcaatcagct tttgtatgac atccgaacta atgcagtcac 1020
cgtgggtggt gtggcagctg gaattggggg cctcgagctg gctgccatga ttgtgtccat 1080
gtatetgtae tgeaatetae aataagteea ettetgeete tgecaetaet getgeeacat 1140
caatgtcact tgggccagaa tggacctgcc ctttctgctc cagacttggg gctagatagg 1260
gaccactect tttaggegat geetgaettt cetteeattg gtgggtggat gggtgggggg 1320
cattccagag cctctaaggt agccagttct gttgcccatt cccccagtct attaaaccct 1380
tgatatgccc cctaggccta gtggtgatcc cagtgctcta ctgggggatg agagaaaggc 1440
attttatagc ctgggcataa gtgaaatcag cagagcctct gggtggatgt gtagaaggca 1500
cttcaaaatg cataaacctg ttacaatgtt gcc
                                                                 1533
<210> 15
<211> 472
<212> DNA
<213> Homo sapiens
<400> 15
tcagagaaaa ctcaaacttt attgagagaa ttttcaaatt ttcagtcaca ttttcaatgt 60
gacatcagec atgtgtgtag cttcagettg tettetttt aacttatgge tgeceatete 120
ctgcttcttt agtcttagca tgcttaggat taggtggagt cttctcttt acatcagagc 180
catctccacg ctcactccga gtcttttcca gatccatttc ctggcaatca ccttctactt 240
tacgttette gateggaggt gtteettete tetettgtee aggtteaata teetgattgt 300
cagttggtgg ttcctcttgc tgagattcac cgggagccac gaatgcaacc acatcgggag 360
ectectgace atetectett cetetggate ttgateteae tegtgeaete ategetgeaa 420
```

```
ctagaagatc gtgaactgaa gaacttgagt cagcagagag cctgqcqaag aa
                                                                    472
 <210> 16
 <211> 478
 <212> DNA
 <213> Homo sapiens
 <400> 16
 cttcattctt cgccaggctc tctgctgact caagttcttc agttcacgat cttctagttg 60
 cagcgatgag tgcacgagtg agatcaagat ccagaggaag aggagatggt caggaggctc 120
 ccgatgtggt tgcattcgtg gctcccggtg aatctcagca agaggaacca ccaactgaca 180
 atcaggatat tgaacctgga caagagagag aaggaacacc tccgatcgaa gaacgtaaag 240
 tagaaggtga ttgccaggaa atggatctgg aaaagactcg gagtgagcgt ggagatggct 300
 ctgatgtaaa agagaagact ccacctaatc ctaagcatgc taagactaaa gaagcaggag 360
 atgggcagcc ataagttaaa aagaagacaa gctgaagcta cacacatggc tgatgtcaca 420
 ttgaaaatgt gactgaaaat ttgaaaattc tctcaataaa gtttgagttt tctctgaa
 <210> 17
 <211> 198
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> unsure
<222> (191)
<400> 17
eccgctgtac caccccagca tgttctgcgc cggcggaggg caagaccaga aggactcctg 60
caacggtgac tetgggggge ceetgatetg caacgggtac ttgcagggec ttgtgtettt 120
cggaaaagcc ccgtgtggcc aagttggcgt gccaggtgtc tacaccaacc tctgcaaatt 180
cactgagtgg nattaagg
<210> 18
<211> 465
<212> DNA
<213> Homo sapiens
<400> 18
tggagatgga gtatgtattt attttacaaa aataaatcac catcttcgga ccatttgtag 60
actggaacat ttcgagcaat gagtgcgcca cacggacgag tgccctggtg actccctgat 120
gttegegtea ceeccaggge cacettggeg eeegcatgag cetegettee cacteeegge 180
ctccaactcc cttccctcgc agccgccatt caccttctgc tgtttatttg tctgcagagc 240
gcctggacac cggaaaaggc gattccctga gcgcctggag ttggagacaa ttcctggttc 300
agaatttaaa catctttcta aggtaagcgc tgctccaaaa ctcttcgccg cgtggggact 360
ttgcaccagg ggcggttggg aaggaagttg gccctccacg ggttcctggg caaccgcggc 420
ctgttgaaaa aaggttctgg gtcaaataat ttaacttcgg aggag
```

<210> 19

PCT/US99/24331 WO 00/23111 <211> 204 <212> DNA <213> Homo sapiens <400> 19 ggcgggaaca ggcggcgctg gacctgtacc cctacgacgc cgggacggac agcggcttca 60 cettetecte ecceaactte gecaceatee egeaggacae ggtgacegag ataaegteet 120 ceteteccag ccaceeggee aacteettet actaeeegeg getgaaggee etgeeteeea 180 tcgccagggt gacactggtg cggc 204 <210> 20 <211> 294 <212> DNA <213> Homo sapiens <220> <221> unsure <222> (287) <400> 20 gagatttctc ttcaatggct tcctgtgagc tagagtttga aaatatctta aaatcttgag 60 ctagagatgg aagtagcttg gacgattttc attatcatgt aaatcgggtc actcaagggg 120 ccaaccacag ctgggagcca ctgctcaggg gaaggttcat atgggacttt ctactgccca 180 aggttctata caggatataa aggtgcctca cagtatagat ctggtagcaa agtaagaaga 240 aacaaacact gatetettte tgecacecet etgaceettt ggaactnete tgac 294 <210> 21 <211> 22 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence: Synthetic <400> 21 atcagaacaa agaggctgtg tc 22 <210> 22 <211> 21 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence:Synthetic <400> 22 atctctaaag ccccaacctt c 21

WO 00/23111	PCT/US99/24331
<210> 23	
<211> 19	_
<212> DNA	
<213> Artificial Sequence	Ch. ··
	•
<220>	
<223> Description of Artificial Sequence:Synthetic	
<400> 23	
tgccgaagag gttcagtgc	19
•	
<210> 24	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:Synthetic	
-400- 74	
<400> 24	22
gccacagtgg tactgtccag at	22
<210> 25	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	•
<223> Description of Artificial Sequence:Synthetic	
<400> 25	
gctgcaagtt ctccacattg a	21
<210> 26	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:Synthetic	
<400> 26	
cagccgcagg tgaaacac	18
- Laguegeagg Egaaacac	10
<210> 27	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
•	

<220>	•	
<223>	Description of Artificial Sequence:Synthetic	
	·	
<400>	27	
tggctt	ttgaa ctcagggtca	20
<210>	28	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Description of Artificial Sequence:Synthetic	
<400>	28	
cggatg	rcacc tegtagacag	20
<210>	29	
<211>	20	
<212>		
<213>	Artificial Sequence	
<220>		
<223>	Description of Artificial Sequence:Synthetic	
	20	
<400>		
eggeaa	cctg gtagtgagtg	20
<210>	30	
<211> 3		
<212> I		
	Artificial Sequence	
	interior dequence	
<220>		
	Description of Artificial Sequence:Synthetic	
	boquonocio, municipal	
<400> 3	30	
cgcagct	cet tgtaaactte ag	22
<210> 3	31	
<211> 2	20	
<212> D	ONA	
<213> A	rtificial Sequence	
	-	
<220>		
<223> D	escription of Artificial Sequence:Synthetic	

WO 00/23111	PCT/US99/24331
<400> 31	
cgggaaccta ccagcctatg	20
<210> 32	& .
<211> 20	•
<212> DNA	
<213> Artificial Sequence	
<220>	·
<223> Description of Artificial Sequence:Synthetic	
<400> 32	
caggcaacag ggagtcatgt	20
<210> 33	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:Synthetic	
<400> 33	
tgggcatctg ggtgtcaa	18
<210> 34	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:Synthetic	
<400> 34	
cggctgcgat gaggaagta	19
<210> 35	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:Synthetic	•
<400> 35	
gcccatctcc tgcttcttta gt	22
<210> 36	

WO 00/23111 PCT/US99/2433

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 36

cgtggagatg gctctgatgt a

21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24331

			· · · · · · · · · · · · · · · · · · ·				
A. CLASSIFICATION OF SUBJECT MATTER							
US CL	IPC(7) :Please See Extra Sheet. US CL :Please See Extra Sheet.						
According	to International Patent Classification (IPC) or to both	h national classification and IPC					
	LDS SEARCHED						
Minimum d	documentation searched (classification system follow	red by classification symbols)					
U.S. ;	424/130.1, 141.1, 155.1, 183.1; 435/6, 7.1, 7.23, 7.	9, 91.2; 436/501, 504, 505, 547; 514/44;	536/23.5				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched							
Electronic o	data base consulted during the international search (name of data base and, where practicable	, search terms used)				
	Biosis, Embase, Cancerlit, Scisearch, WPIDS, USPArms: CSG, cancer specific gene, cancer, diagnosis	ATFULL					
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT						
Category*	Citation of document, with indication, where a	ppropriate, of the relevant passages	Relevant to claim No.				
Y	Database SCISEARCH, Accession Nu Reverse transcriptase-polymerase char cancer. Urologic Clinics of North Am 2, pages 367-&.	in reaction assays for prostate	1-6				
Y	Y CHO-CHUNG et al. Antisense Oligonucleotides for the treatment of cancer. Current Opinion in Therapeutic Patents. 1993, Vol. 3, No. 12, pages 1737-1750, see entire document.						
A,E	BUSSEMAKERS et al. DD3: A new overexpressed in prostate cancer. Ca 1999, Vol. 59, No. 23, pages 5975-5	ancer Research. 01 December	1-7				
Furth	er documents are listed in the continuation of Box C	C. See patent family annex.					
· Spe	scial categories of cited documents:	*T* later document published after the inte	mational filing date or priority				
	cument defining the general state of the art which is not considered be of particular relevance	date and not in conflict with the appli the principle or theory underlying the	cation but cited to understand				
	fier document published on or after the international filing date	"X" document of particular relevance; the					
	cument which may throw doubts on priority claim(s) or which is d to establish the publication date of another citation or other	considered novel or cannot be consider when the document is taken alone					
O doc	cial reason (as specified) cument referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance; the considered to involve an inventive combined with one or more other such	step when the document is documents, such combination				
	ense sument published prior to the international filing date but later than priority date claimed	being obvious to a person skilled in the art *&* document member of the same patent family					
	actual completion of the international search	Date of mailing of the international sear	ch report				
10 FEBRU	10 FEBRUARY 2000 07 MAR 2000						
Commission Box PCT	nailing address of the ISA/US ner of Patents and Trademarks , D.C. 20231	Authorized offices Annae GEETHA P. BANSAL	unce to				
Facsimile No		Telephone No. (703) 308-0196					

INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/24331

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

A61K 39/395, 48/00; C12P 19/34; C12Q 1/68; G01N 33/53, 33/574, 33/546, 33/567

A. CLASSIFICATION OF SUBJECT MATTER: US CL $\,:\,$

424/130.1, 141.1, 155.1, 183.1; 435/6, 7.1, 7.23, 7.9, 91.2; 436/501, 504, 505, 547; 514/44; 536/23.5

Form PCT/ISA/210 (extra sheet)(July 1992)★